arXiv:1206.2557v2 [math.ST] 11 Feb 2013

Asymptotics and multiplier bootstrap of the sequential empirical copula process with applications to change-point detection

Ivan Kojadinovic* and Tom Rohmer
Laboratoire de mathématiques et applications, UMR CNRS 5142
Université de Pau et des Pays de l'Adour
B.P. 1155, 64013 Pau Cedex, France
{ivan.kojadinovic,tom.rohmer}@univ-pau.fr

February 13, 2013

Abstract

The weak limit of the sequential empirical copula process is obtained and the asymptotic validity of a resampling scheme based on multipliers is established under the nonrestrictive smoothness conditions considered by Segers (2012). The empirical process under consideration differs from the sequential process initially studied by Rüschendorf (1976) which cannot be expressed in terms of the empirical copula. The obtained theoretical results are used to derive tests for detecting distributional changes in a sequence of independent continuous observations. The tests are based on the comparison of the empirical copula of the k first and the $n-k$ last observations for all possible values of k. The finite-sample performance of the resulting testing procedures is studied through large-scale Monte Carlo experiments. In the case of distributional changes due to a change in the copula only, the derived tests appear to be substantially more powerful than similar tests recently considered in the literature based on the sequential process initially studied by Rüschendorf.

Keywords: breakpoint detection; empirical copula; multiplier central limit theorem; multivariate independent observations; partial-sum process; ranks.

1 Introduction

Let $\boldsymbol{X}$ be a d-dimensional random vector with continuous marginal cumulative distribution functions (c.d.f.s) $F_1, \dots, F_d$. It is then well-known from the work of Sklar (1959) that the c.d.f. F of $\boldsymbol{X}$ can be written in a unique way as

$$F(\boldsymbol{x}) = C\{F_1(x_1), \dots, F_d(x_d)\}, \qquad \boldsymbol{x} \in \mathbb{R}^d, \tag{1}$$

where the function $C : [0,1]^d \to [0,1]$ is a copula and can be regarded as capturing the dependence between the components of $\boldsymbol{X}$.

Assume that C and $F_1, \dots, F_d$ are unknown, and let $\boldsymbol{X}_i = (X_{i1}, \dots, X_{id})$, $i \in \{1, \dots, n\}$, be a random sample from F. For any $j \in \{1, \dots, d\}$, denote by $R_{ij,n}$ the rank of X_{ij} among $X_{1j}, \dots, X_{nj}$ and let $\hat{U}_{ij,n} = R_{ij,n}/(n+1)$. The random vectors

$$\hat{\boldsymbol{U}}_{i,n} = (\hat{U}_{i1,n}, \dots, \hat{U}_{id,n}), \qquad i \in \{1, \dots, n\}, \tag{2}$$

are frequently referred to as *pseudo-observations* from the copula C.

A natural nonparametric estimator of C is then the *empirical copula* of $\boldsymbol{X}_1, \dots, \boldsymbol{X}_n$ (Rüschendorf, 1976; Deheuvels, 1979, 1981), which is frequently defined as the empirical c.d.f. computed from the pseudo-observations, i.e.,

$$C_n(\boldsymbol{u}) = \frac{1}{n} \sum_{i=1}^{n} \mathbf{1}(\hat{\boldsymbol{U}}_{i,n} \leq \boldsymbol{u}), \qquad \boldsymbol{u} \in [0,1]^d.$$

Note that the quantities $\hat{U}_{ij,n}$ can equivalently be rewritten as $\hat{U}_{ij,n} = n\hat{F}_{n,j}(X_{ij})/(n+1)$, where $\hat{F}_{n,j}$ is the empirical c.d.f. computed from $X_{1j}, \dots, X_{nj}$, and where the scaling factor $n/(n+1)$ is classically introduced to avoid problems at the boundary of $[0,1]^d$.

The object of interest of this paper is the *sequential* empirical copula process defined by

$$\mathbb{C}_n(s, \boldsymbol{u}) = \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} \left\{ \mathbf{1}(\hat{\boldsymbol{U}}_{i,\lfloor ns \rfloor} \leq \boldsymbol{u}) - C(\boldsymbol{u}) \right\}, \qquad (s, \boldsymbol{u}) \in [0,1]^{d+1}, \tag{3}$$

where, for any $y \geq 0$, $\lfloor y \rfloor$ is the greatest integer smaller or equal than y, and, for any $s \in [0,1]$, the $\hat{\boldsymbol{U}}_{i,\lfloor ns \rfloor}$ are pseudo-observations computed analogously to (2) with n replaced by $\lfloor ns \rfloor$. The latter process can be rewritten in terms of the empirical copula $C_{\lfloor ns \rfloor}$ of $\boldsymbol{X}_1, \dots, \boldsymbol{X}_{\lfloor ns \rfloor}$ as

$$\mathbb{C}_n(s, \boldsymbol{u}) = \sqrt{n}\lambda_n(s)\{C_{\lfloor ns \rfloor}(\boldsymbol{u}) - C(\boldsymbol{u})\}, \qquad (s, \boldsymbol{u}) \in [0,1]^{d+1},$$

where $\lambda_n(s) = \lfloor ns \rfloor / n$ and with the convention that $C_0(\boldsymbol{u}) = 0$ for all $\boldsymbol{u} \in [0,1]^d$. For $s = 1$, one recovers the standard empirical copula process which has been extensively studied in the literature (see e.g. Rüschendorf, 1976; Gänssler and Stute, 1987; Fermanian et al., 2004; Tsukahara, 2005; van der Vaart and Wellner, 2007; Segers, 2012; Bücher and Volgushev, 2011). Note that $\mathbb{C}_n$ as defined in (3) differs from the sequential process initially studied by Rüschendorf (1976) and defined by

$$\mathbb{C}_n^R(s, \boldsymbol{u}) = \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} \{\mathbf{1}(\hat{\boldsymbol{U}}_{i,n} \leq \boldsymbol{u}) - C(\boldsymbol{u})\}, \qquad (s, \boldsymbol{u}) \in [0,1]^{d+1}. \tag{4}$$

Unlike $\mathbb{C}_n$, $\mathbb{C}_n^R$ cannot be rewritten in terms of the empirical copula unless $s = 1$. The latter was further studied by Bücher and Volgushev (2011) and was used in Rémillard (2010), Bücher and Ruppert (2013), Wied et al. (2013) and van Kampen and Wied (2012) to derive tests for change-point detection.

The first aim of this paper is to study the asymptotics of the sequential empirical copula process $\mathbb{C}_n$ defined in (3) under the nonrestrictive smoothness conditions considered in Segers (2012), and to show, in this more general framework, the validity of the multiplier bootstrap initially proposed by Rémillard and Scaillet (2009). The second objective of this work is to apply the obtained theoretical results to the derivation of tests for change-point detection for serially independent observations particularly sensitive to changes in the copula.

Given a sequence $\boldsymbol{X}_1, \dots, \boldsymbol{X}_n$ of multivariate independent observations, change-point detection classically aims at testing

$$H_0 : \exists\, F \text{ such that } \boldsymbol{X}_1, \dots, \boldsymbol{X}_n \text{ have c.d.f. } F \tag{5}$$

against alternatives involving the nonconstancy of the c.d.f. Under the assumption that $\boldsymbol{X}_1, \dots, \boldsymbol{X}_n$ have continuous marginal c.d.f.s, from (1), we can write H_0 as $H_{0,m} \cap H_{0,c}$, where

$$H_{0,m} : \exists\, F_1, \dots, F_d \text{ such that } \boldsymbol{X}_1, \dots, \boldsymbol{X}_n \text{ have marginal c.d.f.s } F_1, \dots, F_d, \tag{6}$$

and

$$H_{0,c} : \exists\, C \text{ such that } \boldsymbol{X}_1, \dots, \boldsymbol{X}_n \text{ have copula } C. \tag{7}$$

Following Csörgő and Horváth (1997, Section 2.6) (see also Bai, 1994), a broad class of nonparametric tests for testing H_0 can be derived from the process

$$\mathbb{D}_n^{\#}(s, \boldsymbol{x}) = \sqrt{n}\, \lambda_n(s) \{1 - \lambda_n(s)\} \left\{ F_{\lfloor ns \rfloor}(\boldsymbol{x}) - F^{\star}_{n - \lfloor ns \rfloor}(\boldsymbol{x}) \right\}, \qquad (s, \boldsymbol{x}) \in [0,1] \times \mathbb{R}^d, \tag{8}$$

where

$$F_k(\boldsymbol{x}) = \frac{1}{k} \sum_{i=1}^{k} \mathbf{1}(\boldsymbol{X}_i \leq \boldsymbol{x}) \qquad \text{and} \qquad F^{\star}_{n-k}(\boldsymbol{x}) = \frac{1}{n-k} \sum_{i=k+1}^{n} \mathbf{1}(\boldsymbol{X}_i \leq \boldsymbol{x}), \qquad \boldsymbol{x} \in \mathbb{R}^d,$$

are the empirical c.d.f.s computed from $\boldsymbol{X}_1, \dots, \boldsymbol{X}_k$ and $\boldsymbol{X}_{k+1}, \dots, \boldsymbol{X}_n$, respectively, with the convention that $F_0(\boldsymbol{x}) = 0$ and $F_0^{\star}(\boldsymbol{x}) = 0$ for all $\boldsymbol{x} \in \mathbb{R}^d$. A typical test statistic is then

$$T_n = \max_{1 \leq k \leq n-1} \frac{k(n-k)}{n^{3/2}} \sup_{\boldsymbol{x} \in \mathbb{R}^d} \left| F_k(\boldsymbol{x}) - F^{\star}_{n-k}(\boldsymbol{x}) \right| = \sup_{s \in [0,1]} \sup_{\boldsymbol{x} \in \mathbb{R}^d} |\mathbb{D}_n^{\#}(s, \boldsymbol{x})|,$$

which was for instance used in Gombay and Horváth (1999). While $\mathbb{D}_n^{\#}$ appears as a natural process for testing H_0, typical tests derived from $\mathbb{D}_n^{\#}$ (such as those based on T_n) appear to have little power, in the case of moderate sample size, for detecting distributional changes due to a change in the copula only, i.e., when $H_{0,m} \cap (\neg H_{0,c})$ holds. Empirical evidence of the latter fact can be found in Holmes et al. (2013, Section 4) and will be additionally given in Section 4. For that reason, nonparametric tests for change-point detection particularly sensitive to changes in the dependence structure are of strong practical interest.

Several such tests were proposed in the literature. Tests based on Kendall's tau were investigated by Gombay and Horváth (1999), Gombay and Horváth (2002), Quessy et al. (2013) and Dehling et al. (2013). Tests based on a variant of process (8) related to the process $\mathbb{C}_n^R$ defined in (4) were considered in Rémillard (2010), Bücher and Ruppert (2013), van Kampen and Wied (2012) and Wied et al. (2013). While some of these tests were designed to deal with serially dependent data, to the best of our understanding, none of these tests can be used to directly test $H_{0,c}$ defined in (7) unless $H_{0,m}$ defined in (6) is assumed to hold. In other words, when using these tests, rejection of H_0 should not be interpreted as rejection of $H_{0,c}$ unless one is ready to assume that $H_{0,m}$ holds. This point will be discussed further in Section 5.

The second aim of this work is thus to apply the theoretical results obtained for the process $\mathbb{C}_n$ defined in (3) to derive procedures for testing H_0 based on yet another variant of process (8). As we shall see in Section 3.2, there are theoretical reasons to expect that, at least in some scenarios under $H_{0,m}$, the proposed tests are more powerful than the similar tests considered in Rémillard (2010) and Bücher and Ruppert (2013). The numerical experiments of Section 4 empirically confirm these theoretical hints and reveal that, in finite-samples, the gain in power can be substantial. In a related manner, Corollary 4 in Section 3.2 suggests that, under $H_{0,m}$, should the marginal c.d.f.s $F_1, \dots, F_d$ be known, it is actually preferable to ignore that information, at least in some scenarios, thereby providing yet another illustration of a phenomenon put forward in Genest and Segers (2010) and, more recently, in Bücher (2012).

The paper is organized as follows. In the second section, the weak limit of the sequential empirical copula process $\mathbb{C}_n$ is obtained and the validity of a multiplier bootstrap for $\mathbb{C}_n$ is established under smoothness conditions à la Segers (2012). The third section is devoted to a detailed description of the tests for change-point detection based on the results of Section 2 as well as to a theoretical comparison with the similar tests considered in Rémillard (2010) and Bücher and Ruppert (2013). The fourth section partially reports the results of large-scale Monte Carlo experiments comparing the derived tests with those of Bücher and Ruppert (2013) and a test based on Kendall's tau considered in Quessy et al. (2013), among others. All the proofs are relegated to the appendices.

In the rest of the paper, the arrow '$\rightsquigarrow$' denotes weak convergence in the sense of Definition 1.3.3 in van der Vaart and Wellner (2000), and $\ell^\infty([0,1]^{d+1})$ represents the space of all bounded real-valued functions on $[0,1]^{d+1}$ equipped with the uniform metric.

Note finally that the code of all the tests studied in this work will be released as an R package whose tentative name is `npcp`.

2 The sequential empirical copula process

Let $\boldsymbol{X}_1, \dots, \boldsymbol{X}_n$ be a random sample from c.d.f. F and recall that $F_1, \dots, F_d$ are the continuous univariate margins of F. Then, let $\boldsymbol{U}_i = (U_{i1}, \dots, U_{id})$, $i \in \{1, \dots, n\}$, be the unobservable random sample obtained from $\boldsymbol{X}_1, \dots, \boldsymbol{X}_n$ by the probability integral transforms $U_{ij} = F_j(X_{ij})$, $i \in \{1, \dots, n\}$, $j \in \{1, \dots, d\}$. It follows that $\boldsymbol{U}_1, \dots, \boldsymbol{U}_n$ is a marginally uniform random sample from the unknown c.d.f. C. The corresponding sequential empirical process is then defined as

$$\mathbb{Z}_n(s, \boldsymbol{u}) = \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} \{\mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}) - C(\boldsymbol{u})\}, \qquad (s, \boldsymbol{u}) \in [0,1]^{d+1}. \tag{9}$$

According to Theorem 2.12.1 of van der Vaart and Wellner (2000), $\mathbb{Z}_n \rightsquigarrow \mathbb{Z}_C$ in $\ell^\infty([0,1]^{d+1})$, where $\mathbb{Z}_C$ is a tight centered Gaussian process with covariance function

$$\operatorname{cov}\{\mathbb{Z}_C(s,\boldsymbol{u}), \mathbb{Z}_C(t,\boldsymbol{v})\} = (s \wedge t)\{C(\boldsymbol{u} \wedge \boldsymbol{v}) - C(\boldsymbol{u})C(\boldsymbol{v})\} \tag{10}$$

known as a *C-Kiefer-Müller* process.

To state the weak limit of the sequential empirical copula process $\mathbb{C}_n$ defined in (3), we need the partial derivatives of the copula C. For any $j \in \{1,\dots,d\}$, let $C^{[j]}(\boldsymbol{u})$ be the partial derivative of C with respect to its jth argument at $\boldsymbol{u}$, i.e.,

$$C^{[j]}(\boldsymbol{u}) = \lim_{\substack{h \to 0 \\ u_j + h \in [0,1]}} \frac{C(u_1,\dots,u_{j-1},u_j+h,u_{j+1},\dots,u_d) - C(\boldsymbol{u})}{h}, \qquad \boldsymbol{u} \in [0,1]^d.$$

If $C^{[j]}(\boldsymbol{u})$ exists and is continuous on $[0,1]^d$ for all $j \in \{1,\dots,d\}$, we know from the work of Gänssler and Stute (1987), Fermanian et al. (2004), Tsukahara (2005) or van der Vaart and Wellner (2007) that the empirical copula process $\sqrt{n}(C_n - C) = \mathbb{C}_n(1,\cdot)$ converges weakly in $\ell^\infty([0,1]^d)$ to the tight centered Gaussian process

$$\mathbb{C}_C(1,\boldsymbol{u}) = \mathbb{Z}_C(1,\boldsymbol{u}) - \sum_{j=1}^d C^{[j]}(\boldsymbol{u})\mathbb{Z}_C(1,\boldsymbol{u}^{(j)}), \qquad \boldsymbol{u} \in [0,1]^d, \tag{11}$$

where, for any $j \in \{1,\dots,d\}$ and any $\boldsymbol{u} \in [0,1]^d$, $\boldsymbol{u}^{(j)}$ is the vector of $[0,1]^d$ defined by $u_i^{(j)} = u_j$ if $i = j$ and 1 otherwise, and $\mathbb{Z}_C(1,\cdot)$ is a C-Brownian bridge.

For many copula families however, the partial derivatives $C^{[j]}$, $j \in \{1,\dots,d\}$, fail to be continuous on the whole of $[0,1]^d$. To deal with such situations, Segers (2012) considered the following condition:

($\mathcal{C}1$) for any $j \in \{1,\dots,d\}$, $C^{[j]}$ exists and is continuous on the set $V_j = \{\boldsymbol{u} \in [0,1]^d : 0 < u_j < 1\}$.

Under Condition ($\mathcal{C}1$), for any $j \in \{1,\dots,d\}$, Segers (2012) extended the domain of $C^{[j]}$ to the whole of $[0,1]^d$ by setting

$$C^{[j]}(\boldsymbol{u}) = \begin{cases} \limsup\limits_{h \downarrow 0} \dfrac{C(u_1,\dots,u_{j-1},h,u_{j+1},\dots,u_d)}{h}, & \text{if } \boldsymbol{u} \in [0,1]^d,\ u_j = 0, \\ \limsup\limits_{h \downarrow 0} \dfrac{C(\boldsymbol{u}) - C(u_1,\dots,u_{j-1},1-h,u_{j+1},\dots,u_d)}{h}, & \text{if } \boldsymbol{u} \in [0,1]^d,\ u_j = 1, \end{cases}$$

which ensures that the process $\mathbb{C}_C(1,\cdot)$ defined in (11) is well-defined on the whole of $[0,1]^d$, and showed the weak convergence of the empirical copula process $\sqrt{n}(C_n - C) = \mathbb{C}_n(1,\cdot)$ to $\mathbb{C}_C(1,\cdot)$ in $\ell^\infty([0,1]^d)$. Condition ($\mathcal{C}1$) is nonrestrictive in the sense that it is necessary for $\mathbb{C}_C(1,\cdot)$ to have continuous sample paths and was verified in Segers (2012) for many popular copula families.

A stronger result can be obtained if the following additional smoothness condition is assumed:

($\mathcal{C}2$) for any $i, j \in \{1, \dots, d\}$, the second-order partial derivative $C^{[i,j]} = \partial^2 C/(\partial u_i \partial u_j)$ exists, is continuous on the set $V_i \cap V_j = \{\boldsymbol{u} \in [0,1]^d : u_i, u_j \in (0,1)\}$, and

$$\sup_{\boldsymbol{u} \in V_i \cap V_j} \max\{u_i(1-u_i), u_j(1-u_j)\} |C^{[i,j]}(\boldsymbol{u})| < \infty.$$

Under Conditions ($\mathcal{C}1$) and ($\mathcal{C}2$), Segers (2012, Proposition 4.2) showed that, with probability one,

$$\sup_{\boldsymbol{u} \in [0,1]^d} \left| \mathbb{C}_n(1, \boldsymbol{u}) - \left\{ \mathbb{Z}_n(1, \boldsymbol{u}) - \sum_{j=1}^d C^{[j]}(\boldsymbol{u}) \mathbb{Z}_n(1, \boldsymbol{u}^{(j)}) \right\} \right| = O(n^{-1/4} (\log n)^{1/2} (\log \log n)^{1/4}), \tag{12}$$

thereby recovering the result initially stated by Stute (1984) under very restrictive conditions on C.

The result stated in (12) is at the root of the weak convergence of the sequential empirical copula process $\mathbb{C}_n$ defined in (3). The proof of the following proposition is given in Appendix A.

Proposition 1. *Under Conditions ($\mathcal{C}1$) and ($\mathcal{C}2$), $\mathbb{C}_n \rightsquigarrow \mathbb{C}_C$ in $\ell^\infty([0,1]^{d+1})$, where*

$$\mathbb{C}_C(s, \boldsymbol{u}) = \mathbb{Z}_C(s, \boldsymbol{u}) - \sum_{j=1}^d C^{[j]}(\boldsymbol{u}) \mathbb{Z}_C(s, \boldsymbol{u}^{(j)}), \qquad (s, \boldsymbol{u}) \in [0,1]^{d+1}. \tag{13}$$

The computation of approximate p-values for tests involving the sequential empirical copula process will typically be based on some resampling scheme. In the nonsequential setting, Bücher and Dette (2010) compared the finite-sample behavior of the various resampling techniques proposed in the literature and concluded that the multiplier procedure of Rémillard and Scaillet (2009) has, overall, the best finite-sample properties. This technique was revisited theoretically by Segers (2012) who showed its asymptotic validity under Condition ($\mathcal{C}1$). In the remainder of this section, we show the asymptotic validity of the multiplier approach in the sequential setting under consideration.

Let M be a large integer and let $\xi_i^{(m)}$, $i \in \{1, \dots, n\}$, $m \in \{1, \dots, M\}$, be i.i.d. random variables independent of the data $\boldsymbol{X}_1, \dots, \boldsymbol{X}_n$. We consider two possibilities for the $\xi_i^{(m)}$:

($\mathcal{M}1$) $\xi_i^{(m)}$ has mean 0, variance 1 and satisfies $\int_0^\infty \{\mathrm{P}(|\xi_i^{(m)}| > x)\}^{1/2} \mathrm{d}x < \infty$;

($\mathcal{M}2$) $\xi_i^{(m)}$ is a Rademacher random variable, i.e., $\mathrm{P}(\xi_i^{(m)} = 1) = \mathrm{P}(\xi_i^{(m)} = -1) = 1/2$.

Clearly, a result that holds under ($\mathcal{M}1$) will also hold under ($\mathcal{M}2$).

Next, for any $m \in \{1, \dots, M\}$, let

$$\mathbb{Z}_n^{(m)}(s, \boldsymbol{u}) = \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} \xi_i^{(m)} \{\mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}) - C(\boldsymbol{u})\}, \qquad (s, \boldsymbol{u}) \in [0,1]^{d+1}. \tag{14}$$

A consequence of the sequential extension of the multiplier central limit theorem proved in Holmes et al. (2013, Theorem 1) is that

$$\left(\mathbb{Z}_n, \mathbb{Z}_n^{(1)}, \dots, \mathbb{Z}_n^{(M)}\right) \rightsquigarrow \left(\mathbb{Z}_C, \mathbb{Z}_C^{(1)}, \dots, \mathbb{Z}_C^{(M)}\right)$$

in $\{\ell^\infty([0,1]^{d+1})\}^{M+1}$, where $\mathbb{Z}_C$, a *C-Kiefer-Müller* process, is the weak limit of the sequential empirical process $\mathbb{Z}_n$ defined in (9), and $\mathbb{Z}_C^{(1)}, \dots, \mathbb{Z}_C^{(M)}$ are independent copies of $\mathbb{Z}_C$. For large n, $\mathbb{Z}_n^{(1)}, \dots, \mathbb{Z}_n^{(M)}$ could therefore be regarded as "almost" independent copies of $\mathbb{Z}_n$. Unfortunately, the $\mathbb{Z}_n^{(m)}$ cannot be computed because the random sample $\boldsymbol{U}_1, \dots, \boldsymbol{U}_n$ is unobservable and C is unknown.

We consider two versions of the $\mathbb{Z}_n^{(m)}$ that can be computed depending on how the $\boldsymbol{U}_i$ and C are estimated. For any $s \in [0,1]$ and $\boldsymbol{u} \in [0,1]^d$, let

$$\hat{\mathbb{Z}}_n^{(m)}(s, \boldsymbol{u}) = \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} \xi_i^{(m)} \left\{ \mathbf{1}(\hat{\boldsymbol{U}}_{i,\lfloor ns \rfloor} \leq \boldsymbol{u}) - C_{\lfloor ns \rfloor}(\boldsymbol{u}) \right\} = \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} (\xi_i^{(m)} - \bar{\xi}_{\lfloor ns \rfloor}^{(m)}) \mathbf{1}(\hat{\boldsymbol{U}}_{i,\lfloor ns \rfloor} \leq \boldsymbol{u}), \tag{15}$$

where the $\hat{\boldsymbol{U}}_{i,\lfloor ns \rfloor}$ are defined as in (2) with n replaced by $\lfloor ns \rfloor$, $\bar{\xi}_{\lfloor ns \rfloor}^{(m)} = \lfloor ns \rfloor^{-1} \sum_{i=1}^{\lfloor ns \rfloor} \xi_i^{(m)}$ and $\bar{\xi}_0^{(m)} = 0$ by convention, and let

$$\check{\mathbb{Z}}_n^{(m)}(s, \boldsymbol{u}) = \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} \xi_i^{(m)} \left\{ \mathbf{1}(\hat{\boldsymbol{U}}_{i,n} \leq \boldsymbol{u}) - C_n(\boldsymbol{u}) \right\}. \tag{16}$$

To define "almost" independent copies of $\mathbb{C}_n$ for large n in the spirit of Rémillard and Scaillet (2009), we also need to estimate the partial derivatives $C^{[j]}$ appearing in (13). To do so, we consider the estimator $C_n^{[j]}$ defined in Kojadinovic et al. (2011) by

$$\begin{aligned} C_n^{[j]}(\boldsymbol{u}) = \frac{1}{u_{j,n}^+ - u_{j,n}^-} \{ & C_n(u_1, \dots, u_{j-1}, u_{j,n}^+, u_{j+1}, \dots, u_d) \\ & - C_n(u_1, \dots, u_{j-1}, u_{j,n}^-, u_{j+1}, \dots, u_d) \}, \qquad \boldsymbol{u} \in [0,1]^d, \end{aligned} \tag{17}$$

where $u_{j,n}^+ = (u_j + n^{-1/2}) \wedge 1$, and $u_{j,n}^- = (u_j - n^{-1/2}) \vee 0$. This estimator differs slightly from the one initially proposed in Rémillard and Scaillet (2009). It has the advantage of converging in probability to $C^{[j]}$ uniformly over $[0,1]^d$ if $C^{[j]}$ happens to be continuous on $[0,1]^d$ instead of only satisfying Condition $(\mathcal{C}1)$.

The following result, proved in Appendix A, complements Lemma 2 of Kojadinovic et al. (2011).

Proposition 2. *Let $0 < a < b < 1$ and assume that Conditions $(\mathcal{C}1)$ and $(\mathcal{C}2)$ hold. Then, for any $j \in \{1, \dots, d\}$,*

$$\sup_{\substack{\boldsymbol{u} \in [0,1]^d \\ u_j \in [a,b]}} |C_n^{[j]}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{u})| = O\left(n^{-1/4} (\log n)^{1/2} (\log \log n)^{1/4}\right), \qquad \textit{almost surely.}$$

We can now define empirical processes that can be fully computed and that, under appropriate conditions, can be regarded as “almost” independent copies of $\mathbb{C}_n$ for large n. For any $m \in \{1, \dots, M\}$, $s \in [0, 1]$ and $\boldsymbol{u} \in [0, 1]^d$, let

$$\hat{\mathbb{C}}_n^{(m)}(s, \boldsymbol{u}) = \hat{\mathbb{Z}}_n^{(m)}(s, \boldsymbol{u}) - \sum_{j=1}^{d} C_{\lfloor ns \rfloor}^{[j]}(\boldsymbol{u}) \hat{\mathbb{Z}}_n^{(m)}(s, \boldsymbol{u}^{(j)}) \tag{18}$$

and

$$\check{\mathbb{C}}_n^{(m)}(s, \boldsymbol{u}) = \check{\mathbb{Z}}_n^{(m)}(s, \boldsymbol{u}) - \sum_{j=1}^{d} C_n^{[j]}(\boldsymbol{u}) \check{\mathbb{Z}}_n^{(m)}(s, \boldsymbol{u}^{(j)}). \tag{19}$$

The following result, proved in Appendix B, can be seen as an extension of Proposition 3.2 of Segers (2012) to the sequential setting.

Theorem 1. *Assume that Conditions ($\mathcal{C}1$) and ($\mathcal{C}2$) hold.*

(i) Under ($\mathcal{M}1$),

$$\left(\mathbb{C}_n, \check{\mathbb{C}}_n^{(1)}, \dots, \check{\mathbb{C}}_n^{(M)}\right) \rightsquigarrow \left(\mathbb{C}_C, \mathbb{C}_C^{(1)}, \dots, \mathbb{C}_C^{(M)}\right),$$

in $\{\ell^\infty([0, 1]^{(d+1)})\}^{M+1}$, where $\mathbb{C}_C$ is the weak limit of $\mathbb{C}_n$ and $\mathbb{C}_C^{(1)}, \dots, \mathbb{C}_C^{(M)}$ are independent copies $\mathbb{C}_C$.

(ii) If ($\mathcal{M}2$) is assumed instead of ($\mathcal{M}1$),

$$\left(\mathbb{C}_n, \hat{\mathbb{C}}_n^{(1)}, \dots, \hat{\mathbb{C}}_n^{(M)}, \check{\mathbb{C}}_n^{(1)}, \dots, \check{\mathbb{C}}_n^{(M)}\right) \rightsquigarrow \left(\mathbb{C}_C, \mathbb{C}_C^{(1)}, \dots, \mathbb{C}_C^{(M)}, \mathbb{C}_C^{(1)}, \dots, \mathbb{C}_C^{(M)}\right),$$

in $\{\ell^\infty([0, 1]^{(d+1)})\}^{2M+1}$.

3 Tests for a change in the dependence structure

The setting is as follows. Given a sequence of independent continuous d-dimensional random vectors $\boldsymbol{X}_1, \dots, \boldsymbol{X}_n$, the objective is to test H_0 defined in (5) against alternatives involving the nonconstancy of the c.d.f. The aim being to derive tests particularly sensitive to changes in the copula of the sequence, the behavior of the proposed tests will be theoretically investigated under the alternative hypothesis of constant marginal c.d.f.s and a single change-point in the copula, i.e., under $H_{0,m} \cap H_{1,c}$, where $H_{0,m}$ is defined in (6) and $H_{1,c}$ is given by:

$$\begin{aligned} H_{1,c} : \exists \text{ distinct } C_1 \text{ and } C_2, \text{ and } k^\star \in \{1, \dots, n-1\} \text{ such that} \\ \boldsymbol{X}_1, \dots, \boldsymbol{X}_{k^\star} \text{ have copula } C_1 \text{ and } \boldsymbol{X}_{k^\star+1}, \dots, \boldsymbol{X}_n \text{ have copula } C_2. \end{aligned} \tag{20}$$

After describing the proposed tests, we will compare them with the similar tests considered in Rémillard (2010, Section 5.2) and Bücher and Ruppert (2013, Section 3.2), and provide theoretical hints suggesting that, under $H_{0,m}$, the former should be more powerful than the latter at least in some scenarios.

3.1 Tests based on the empirical copula

As explained in the introduction, the aim is to adapt to the current copula context the approach studied in detail in Csörgő and Horváth (1997, Section 2.6) based on the process defined in (8). The idea consists of comparing, for all $k \in \{1, \dots, n-1\}$, the empirical copula of $\boldsymbol{X}_1, \dots, \boldsymbol{X}_k$ with the empirical copula of $\boldsymbol{X}_{k+1}, \dots, \boldsymbol{X}_n$, respectively defined by

$$C_k(\boldsymbol{u}) = \frac{1}{k} \sum_{i=1}^{k} \mathbf{1}(\hat{\boldsymbol{U}}_{i,k} \leq \boldsymbol{u}) \qquad \text{and} \qquad C^\star_{n-k}(\boldsymbol{u}) = \frac{1}{n-k} \sum_{i=k+1}^{n} \mathbf{1}(\hat{\boldsymbol{U}}^\star_{i,n-k} \leq \boldsymbol{u}), \quad \boldsymbol{u} \in [0,1]^d,$$

where $\hat{\boldsymbol{U}}^\star_{k+1,n-k}, \dots, \hat{\boldsymbol{U}}^\star_{n,n-k}$ are the pseudo-observations computed from $\boldsymbol{X}_{k+1}, \dots, \boldsymbol{X}_n$ similarly to (2). Analogous to (8), we define the process

$$\mathbb{D}_n(s, \boldsymbol{u}) = \sqrt{n}\, \lambda_n(s) \{1 - \lambda_n(s)\} \{C_{\lfloor ns \rfloor}(\boldsymbol{u}) - C^\star_{n-\lfloor ns \rfloor}(\boldsymbol{u})\}, \qquad (s, \boldsymbol{u}) \in [0,1]^{d+1}, \quad (21)$$

where $\lambda_n(s) = \lfloor ns \rfloor / n$ and with the convention that $C_0(\boldsymbol{u}) = 0$ and $C^\star_0(\boldsymbol{u}) = 0$ for all $\boldsymbol{u} \in [0,1]^d$.

The weak limit of $\mathbb{D}_n$ is established in the following result proved in Appendix C.

Proposition 3. *Under Conditions (*$\mathcal{C}1$*) and (*$\mathcal{C}2$*), and* H_0*,* $\mathbb{D}_n \rightsquigarrow \mathbb{D}_C$ *in* $\ell^\infty([0,1]^{d+1})$*, where, for any* $(s, \boldsymbol{u}) \in [0,1]^{d+1}$*,* $\mathbb{D}_C(s, \boldsymbol{u}) = \mathbb{C}_C(s, \boldsymbol{u}) - s\mathbb{C}_C(1, \boldsymbol{u})$*, with* $\mathbb{C}_C$ *defined in (13), and where the covariance function of* $\mathbb{D}_C$ *can be expressed in terms of the covariance function of* $\mathbb{C}_C$ *as*

$$\operatorname{cov}\{\mathbb{D}_C(s, \boldsymbol{u}), \mathbb{D}_C(t, \boldsymbol{v})\} = (s \wedge t - st) \operatorname{cov}\{\mathbb{C}_C(1, \boldsymbol{u}), \mathbb{C}_C(1, \boldsymbol{v})\}. \quad (22)$$

To compute approximate p-values for tests based on $\mathbb{D}_n$, we adapt the multiplier approach described in the previous section to the current change-point setting. Analogous to $\hat{\mathbb{Z}}_n^{(m)}$ and $\check{\mathbb{Z}}_n^{(m)}$ defined in (15) and (16), respectively, for any $m \in \{1, \dots, M\}$, $s \in [0,1]$ and $\boldsymbol{u} \in [0,1]^d$, let

$$\begin{aligned} \hat{\mathbb{Z}}_n^{\star,(m)}(s, \boldsymbol{u}) &= \frac{1}{\sqrt{n}} \sum_{i=\lfloor ns \rfloor + 1}^{n} \xi_i^{(m)} \left\{ \mathbf{1}(\hat{\boldsymbol{U}}^\star_{i,n-\lfloor ns \rfloor} \leq \boldsymbol{u}) - C^\star_{n-\lfloor ns \rfloor}(\boldsymbol{u}) \right\} \\ &= \frac{1}{\sqrt{n}} \sum_{i=\lfloor ns \rfloor + 1}^{n} (\xi_i^{(m)} - \bar{\xi}^{\star,(m)}_{n-\lfloor ns \rfloor}) \mathbf{1}(\hat{\boldsymbol{U}}^\star_{i,n-\lfloor ns \rfloor} \leq \boldsymbol{u}), \end{aligned}$$

where $\bar{\xi}^{\star,(m)}_{n-\lfloor ns \rfloor} = (n - \lfloor ns \rfloor)^{-1} \sum_{i=\lfloor ns \rfloor+1}^{n} \xi_i^{(m)}$ and $\bar{\xi}^{\star,(m)}_0 = 0$ by convention, and let

$$\check{\mathbb{Z}}_n^{\star,(m)}(s, \boldsymbol{u}) = \frac{1}{\sqrt{n}} \sum_{i=\lfloor ns \rfloor + 1}^{n} \xi_i^{(m)} \left\{ \mathbf{1}(\hat{\boldsymbol{U}}_{i,n} \leq \boldsymbol{u}) - C_n(\boldsymbol{u}) \right\} = \check{\mathbb{Z}}_n^{(m)}(1, \boldsymbol{u}) - \check{\mathbb{Z}}_n^{(m)}(s, \boldsymbol{u}).$$

Then, for any $s \in [0,1]$ and $\boldsymbol{u} \in [0,1]^d$, let

$$\hat{\mathbb{C}}_n^{\star,(m)}(s, \boldsymbol{u}) = \hat{\mathbb{Z}}_n^{\star,(m)}(s, \boldsymbol{u}) - \sum_{j=1}^{d} C^{[j],\star}_{n-\lfloor ns \rfloor}(\boldsymbol{u}) \hat{\mathbb{Z}}_n^{\star,(m)}(s, \boldsymbol{u}^{(j)}),$$

where $C^{[j],\star}_{n-\lfloor ns \rfloor}$ is the version of the estimator of $C^{[j]}$ defined in (17) computed from $\boldsymbol{X}_{\lfloor ns \rfloor+1}, \dots, \boldsymbol{X}_n$, and let

$$\check{\mathbb{C}}_n^{\star,(m)}(s,\boldsymbol{u}) = \check{\mathbb{Z}}_n^{\star,(m)}(s,\boldsymbol{u}) - \sum_{j=1}^d C_n^{[j]}(\boldsymbol{u}) \check{\mathbb{Z}}_n^{\star,(m)}(s,\boldsymbol{u}^{(j)}).$$

Finally, for any $m \in \{1,\dots,M\}$, $s \in [0,1]$ and $\boldsymbol{u} \in [0,1]^d$, we form

$$\hat{\mathbb{D}}_n^{(m)}(s,\boldsymbol{u}) = \{1-\lambda_n(s)\}\, \hat{\mathbb{C}}_n^{(m)}(s,\boldsymbol{u}) - \lambda_n(s) \hat{\mathbb{C}}_n^{\star,(m)}(s,\boldsymbol{u}),$$

and

$$\check{\mathbb{D}}_n^{(m)}(s,\boldsymbol{u}) = \{1-\lambda_n(s)\}\, \check{\mathbb{C}}_n^{(m)}(s,\boldsymbol{u}) - \lambda_n(s) \check{\mathbb{C}}_n^{\star,(m)}(s,\boldsymbol{u}) = \check{\mathbb{C}}_n^{\star,(m)}(s,\boldsymbol{u}) - \lambda_n(s) \check{\mathbb{C}}_n^{\star,(m)}(1,\boldsymbol{u}),$$

where $\hat{\mathbb{C}}_n^{(m)}$ and $\check{\mathbb{C}}_n^{(m)}$ are defined in (18) and (19), respectively.

The following result, proved in Appendix C, is a consequence of Theorem 1.

Proposition 4. *Assume that Conditions ($\mathcal{C}1$) and ($\mathcal{C}2$) hold.*

(i) Under ($\mathcal{M}1$) and H_0,

$$\left(\mathbb{D}_n, \check{\mathbb{D}}_n^{(1)}, \dots, \check{\mathbb{D}}_n^{(M)}\right) \rightsquigarrow \left(\mathbb{D}_C, \mathbb{D}_C^{(1)}, \dots, \mathbb{D}_C^{(M)}\right),$$

in $\{\ell^\infty([0,1]^{(d+1)})\}^{M+1}$, where $\mathbb{D}_C$ is the weak limit of $\mathbb{D}_n$ and $\mathbb{D}_C^{(1)}, \dots, \mathbb{D}_C^{(M)}$ are independent copies $\mathbb{D}_C$.

(ii) If ($\mathcal{M}2$) is assumed instead of ($\mathcal{M}1$), then, under H_0,

$$\left(\mathbb{D}_n, \hat{\mathbb{D}}_n^{(1)}, \dots, \hat{\mathbb{D}}_n^{(M)}, \check{\mathbb{D}}_n^{(1)}, \dots, \check{\mathbb{D}}_n^{(M)}\right) \rightsquigarrow \left(\mathbb{D}_C, \mathbb{D}_C^{(1)}, \dots, \mathbb{D}_C^{(M)}, \mathbb{D}_C^{(1)}, \dots, \mathbb{D}_C^{(M)}\right),$$

in $\{\ell^\infty([0,1]^{(d+1)})\}^{2M+1}$.

The next result, also proved in Appendix C, will be useful to derive the asymptotic behavior of tests based on $\mathbb{D}_n$ under $H_{0,m} \cap H_{1,c}$, where $H_{0,m}$ and $H_{1,c}$ are defined in (6) and (20), respectively.

Proposition 5. *Assume that $H_{0,m} \cap H_{1,c}$ holds with $k^\star = \lfloor nt \rfloor$ for some $t \in (0,1)$. Then,*

(i) $n^{-1/2}\mathbb{D}_n \xrightarrow{\mathrm{P}} K_t$ in $\ell^\infty([0,1]^{d+1})$, where $K_t(s,\boldsymbol{u}) = (s \wedge t)(1 - s \vee t)\{C_1(\boldsymbol{u}) - C_2(\boldsymbol{u})\}$,

(ii) $\sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} |\hat{\mathbb{D}}_n^{(m)}(s,\boldsymbol{u})| = O_{\mathrm{P}}(1)$ and $\sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} |\check{\mathbb{D}}_n^{(m)}(s,\boldsymbol{u})| = O_{\mathrm{P}}(1)$ for all $m \in \{1,\dots,M\}$.

In the framework under consideration, a change in the c.d.f. of the sequence $\boldsymbol{X}_1, \dots, \boldsymbol{X}_n$ can occur at any point $k \in \{1,\dots,n-1\}$. As classically done, a test for change-point detection can therefore be obtained by first defining a test statistic for any possible change-point $k \in \{1,\dots,n-1\}$, and then by combining the resulting $n-1$ statistics into a global statistic.

Natural choices for the $n-1$ change-point statistics are Cramér–von Mises, Kolmogorov-Smirnov and Kuiper statistics (see e.g. Horváth and Shao, 2007; Bücher and Ruppert, 2013). As the latter two statistics led to consistently less powerful tests in our Monte Carlo experiments, for the sake of brevity, we only describe the tests based on the $n-1$ Cramér–von Mises change-point statistics defined by

$$S_{n,k} = \int_{[0,1]^d} \left\{ \mathbb{D}_n \left(\frac{k}{n}, \boldsymbol{u} \right) \right\}^2 \mathrm{d}C_n(\boldsymbol{u}), \qquad k \in \{1,\dots,n-1\}. \tag{23}$$

A natural way of combining $S_{n,1},\dots,S_{n,n-1}$ consists of taking their maximum. This leads to the maximally selected Cramér–von Mises statistic

$$S_n = \max_{1\le k\le n-1} S_{n,k} = \sup_{s\in[0,1]} \int_{[0,1]^d} \{\mathbb{D}_n(s,\boldsymbol{u})\}^2 \mathrm{d}C_n(\boldsymbol{u}). \tag{24}$$

Note that we did also consider the arithmetic mean as combining function as in Holmes et al. (2013) but do not insist on this point further as the resulting tests did not appear more powerful than those based on S_n in our Monte Carlo experiments.

We studied two versions of the test based on S_n depending on whether the approximate p-value is computed using the multiplier processes $\hat{\mathbb{D}}_n^{(m)}$ or the processes $\check{\mathbb{D}}_n^{(m)}$. For any $m \in \{1,\dots,M\}$, let

$$\hat{S}_n^{(m)} = \sup_{s\in[0,1]} \int_{[0,1]^d} \left\{ \hat{\mathbb{D}}_n^{(m)}(s,\boldsymbol{u}) \right\}^2 \mathrm{d}C_n(\boldsymbol{u}) \quad \text{and} \quad \check{S}_n^{(m)} = \sup_{s\in[0,1]} \int_{[0,1]^d} \left\{ \check{\mathbb{D}}_n^{(m)}(s,\boldsymbol{u}) \right\}^2 \mathrm{d}C_n(\boldsymbol{u}).$$

The following result is then a corollary of Proposition 4 and can be proved along the lines of Proposition 1 in Holmes et al. (2013).

Corollary 1. *Assume that Conditions ($\mathcal{C}1$) and ($\mathcal{C}2$) hold.*

(i) Under ($\mathcal{M}1$) and H_0,

$$\left(S_n, \check{S}_n^{(1)}, \dots, \check{S}_n^{(M)} \right) \rightsquigarrow \left(S, S^{(1)}, \dots, S^{(M)} \right)$$

in $[0,\infty)^{(M+1)}$, where

$$S = \sup_{s\in[0,1]} \int_{[0,1]^d} \{\mathbb{D}_C(s,\boldsymbol{u})\}^2 \mathrm{d}C(\boldsymbol{u})$$

is the weak limit of S_n, and $S^{(1)},\dots,S^{(M)}$ are independent copies of S.

(ii) If ($\mathcal{M}2$) is assumed instead of ($\mathcal{M}1$), then, under H_0,

$$\left(S_n, \hat{S}_n^{(1)}, \dots, \hat{S}_n^{(M)}, \check{S}_n^{(1)}, \dots, \check{S}_n^{(M)} \right) \rightsquigarrow \left(S, S^{(1)}, \dots, S^{(M)}, S^{(1)}, \dots, S^{(M)} \right)$$

in $[0,\infty)^{(2M+1)}$.

The previous corollary suggests interpreting the $\hat{S}_n^{(m)}$ (resp. the $\check{S}_n^{(m)}$) under the null hypothesis as M "almost" independent copies of S_n and thus computing an approximate p-value for S_n as

$$\frac{1}{M}\sum_{m=1}^{M}\mathbf{1}\left(\hat{S}_n^{(m)} \geq S_n\right) \qquad \text{or as} \qquad \frac{1}{M}\sum_{m=1}^{M}\mathbf{1}\left(\check{S}_n^{(m)} \geq S_n\right). \tag{25}$$

The following result is a corollary of Proposition 5 and can be proved along the lines of Proposition 2 in Holmes et al. (2013).

Corollary 2. *Assume that $H_{0,m} \cap H_{1,c}$ holds with $k^\star = \lfloor nt \rfloor$ for some $t \in (0,1)$. Then,*

(i) $s \mapsto \int_{[0,1]^d} \frac{\{\mathbb{D}_n(s,\boldsymbol{u})\}^2}{n} \mathrm{d}C_n(\boldsymbol{u}) \quad \overset{\mathrm{P}}{\to} \quad s \mapsto \int_{[0,1]^d} \{K_t(s,\boldsymbol{u})\}^2 \mathrm{d}\{tC_1(\boldsymbol{u}) + (1-t)C_2(\boldsymbol{u})\}$ *in $\ell^\infty([0,1])$, which implies that $S_n \overset{\mathrm{P}}{\to} +\infty$,*

(ii) for any $m \in \{1,\dots,M\}$, $\hat{S}_n^{(m)} = O_{\mathrm{P}}(1)$ and $\check{S}_n^{(m)} = O_{\mathrm{P}}(1)$.

A consequence of the previous corollary is that, under $H_{0,m} \cap H_{1,c}$, the approximate p-values for S_n defined in (25) will tend to zero in probability.

Finally, should H_0 be rejected, and assuming that $H_{0,m} \cap H_{1,c}$ holds with $k^\star = \lfloor nt \rfloor$ for some $t \in (0,1)$, it is frequently of interest in applications to estimate t. In the setting under consideration, an estimator of t can be defined as $t_n = n^{-1} \operatorname{argmax}_{1 \leq k \leq n-1} S_{n,k}$, where the $S_{n,k}$ are defined in (23).

Corollary 3. *Assume that $H_{0,m} \cap H_{1,c}$ holds with $k^\star = \lfloor nt \rfloor$ for some $t \in (0,1)$. Then, $t_n \overset{\mathrm{P}}{\to} t$.*

The above result follows from Corollary 2 (i), the fact that, as a function on $[0,1]$, $s \mapsto \int_{[0,1]^d}\{K_t(s,\boldsymbol{u})\}^2 \mathrm{d}\{tC_1(\boldsymbol{u}) + (1-t)C_2(\boldsymbol{u})\}$ attains its unique maximum at $s = t$, and the argmax theorem (van der Vaart and Wellner, 2000, Corollary 3.2.3); see also Bücher and Ruppert (2013, Proof of Proposition 3) and Dehling et al. (2013, Proof of Theorem 4.3).

3.2 Comparison with tests related to the sequential empirical process of Rüschendorf

Similar tests to those derived in the previous subsection were proposed by Rémillard (2010, Section 5.2) and Bücher and Ruppert (2013, Section 3.2) using the empirical process

$$\mathbb{D}_n^R(s,\boldsymbol{u}) = \sqrt{n}\,\lambda_n(s)\,\{1-\lambda_n(s)\}\,\{\tilde{C}_{\lfloor ns \rfloor}(\boldsymbol{u}) - \tilde{C}^\star_{n-\lfloor ns \rfloor}(\boldsymbol{u})\}, \qquad (s,\boldsymbol{u}) \in [0,1]^{d+1}, \tag{26}$$

where, for any $k \in \{1,\dots,n-1\}$, $\tilde{C}_k$ and $\tilde{C}^\star_{n-k}$ denote respectively the empirical c.d.f.s of the k first and $n-k$ last pseudo-observations $\hat{\boldsymbol{U}}_{1,n},\dots,\hat{\boldsymbol{U}}_{n,n}$ defined in (2), i.e.,

$$\tilde{C}_k(\boldsymbol{u}) = \frac{1}{k}\sum_{i=1}^{k}\mathbf{1}(\hat{\boldsymbol{U}}_{i,n} \leq \boldsymbol{u}) \qquad \text{and} \qquad \tilde{C}^\star_{n-k}(\boldsymbol{u}) = \frac{1}{n-k}\sum_{i=k+1}^{n}\mathbf{1}(\hat{\boldsymbol{U}}_{i,n} \leq \boldsymbol{u}), \quad \boldsymbol{u} \in [0,1]^d,$$

and with the convention that $\tilde{C}_0(\boldsymbol{u}) = 0$ and $\tilde{C}_0^\star(\boldsymbol{u}) = 0$. Starting from $\mathbb{D}_n^R$, these authors naturally suggested to consider maximally selected Kolmogorov–Smirnov, Kuiper and Cramér–von Mises test statistics. The latter is given by

$$S_n^R = \sup_{s \in [0,1]} \int_{[0,1]^d} \left\{ \mathbb{D}_n^R(s, \boldsymbol{u}) \right\}^2 \mathrm{d}C_n(\boldsymbol{u}), \tag{27}$$

and is the analogue of the statistic S_n defined in (24) from the process $\mathbb{D}_n$ defined in (21).

Actually, all the test statistics considered in Rémillard (2010, Section 5.2) and Bücher and Ruppert (2013, Section 3.2) are rank-based, and, consequently, can be rewritten in terms of the process $\mathbb{D}_n^\#$ defined in (8). For instance, in the case of S_n^R, one obtains

$$S_n^R = \sup_{s \in [0,1]} \int_{\mathbb{R}^d} \left\{ \mathbb{D}_n^\#(s, \boldsymbol{x}) \right\}^2 \mathrm{d}F_n(\boldsymbol{x}), \tag{28}$$

where F_n is the empirical c.d.f. computed from $\boldsymbol{X}_1, \dots, \boldsymbol{X}_n$. It follows that the corresponding tests should not be regarded as tests for change-point detection focusing in particular on changes in the copula, but rather as general-purpose tests for detecting changes in the multivariate c.d.f.

The finite-sample performance of the test based on S_n^R (seen as a functional of $\mathbb{D}_n^\#$) was actually also investigated in Holmes et al. (2013, Section 4). The test was found to have good power for detecting changes in the marginal c.d.f.s. but did not appear to detect changes in the copula well. Similar empirical findings will be presented in the next section.

Let us now assume that $H_{0,m}$ defined in (6) holds. In the remainder of this section, we provide theoretical hints suggesting that, at least in some situations, tests based on $\mathbb{D}_n$ should be more powerful than analogous tests based on $\mathbb{D}_n^R$. The main ingredient of our argumentation is a covariance inequality obtained by Genest and Segers (2010).

From Theorem 5 of Bücher and Ruppert (2013), we have that, under H_0, $\mathbb{D}_n^R \rightsquigarrow \mathbb{D}_C^R$ in $\ell^\infty([0,1]^{d+1})$, where $\mathbb{D}_C^R(s, \boldsymbol{u}) = \mathbb{Z}_C(s, \boldsymbol{u}) - s\mathbb{Z}_C(1, \boldsymbol{u})$ for all $(s, \boldsymbol{u}) \in [0,1]^{d+1}$, and $\mathbb{Z}_C$ is a C-Kiefer-Müller process whose covariance function is given in (10). Simple calculations show that the covariance function of $\mathbb{D}_C^R$ can be expressed in terms of the covariance function of the C-Brownian bridge $\mathbb{Z}_C(1, \cdot)$ as

$$\operatorname{cov}\{\mathbb{D}_C^R(s, \boldsymbol{u}), \mathbb{D}_C^R(t, \boldsymbol{v})\} = (s \wedge t - st) \operatorname{cov}\{\mathbb{Z}_C(1, \boldsymbol{u}), \mathbb{Z}_C(1, \boldsymbol{v})\}.$$

Combining the above expression, (22) and Proposition 1 of Genest and Segers (2010), we obtain the following result.

Corollary 4. *Assume that Condition ($\mathcal{C}1$) holds and that C is a bivariate copula that is left-tail decreasing in both arguments. Then, under H_0, for any $(s, \boldsymbol{u})$, $(t, \boldsymbol{v}) \in [0,1]^3$,*

$$\operatorname{cov}\{\mathbb{D}_C(s, \boldsymbol{u}), \mathbb{D}_C(t, \boldsymbol{v})\} \leq \operatorname{cov}\{\mathbb{D}_C^R(s, \boldsymbol{u}), \mathbb{D}_C^R(t, \boldsymbol{v})\}. \tag{29}$$

From Nelsen (2006, Section 5.2.2 and Exercise 5.35), a bivariate copula C is left-tail decreasing (LTD) in both arguments if, for any $0 < u \leq u' \leq 1$ and $0 < v \leq v' \leq 1$, $C(u, v)/(uv) \geq C(u', v')/(u'v')$. As shown by Garralda-Guillem (2000), extreme-value copulas are LTD in both arguments but so are the most popular bivariate copulas with positive dependence such as the Clayton, Frank, normal, t and Plackett.

To illustrate a possible consequence of (29) and provide some intuition, assume that $d = 2$, that Condition ($\mathcal{C}1$) holds and that C is LTD in both arguments. Furthermore, suppose that test statistics are defined using the map $\mathcal{T}$ from $\ell^\infty([0,1]^3)$ to $\mathbb{R}$ given by $\mathcal{T}(D) = \int_{[0,1]^3} D(s, \boldsymbol{u}) \mathrm{d}s \mathrm{d}\boldsymbol{u}$, $D \in \ell^\infty([0,1]^3)$, and that large values of $|\mathcal{T}(\mathbb{D}_n^R)|$ and $|\mathcal{T}(\mathbb{D}_n)|$ lead to the rejection of H_0. Then, from the continuous mapping theorem and (29), we obtain that, under H_0, $\mathcal{T}(\mathbb{D}_n^R) \rightsquigarrow \mathcal{T}(\mathbb{D}_C^R)$ and $\mathcal{T}(\mathbb{D}_n) \rightsquigarrow \mathcal{T}(\mathbb{D}_C)$, where $\mathcal{T}(\mathbb{D}_C^R)$ and $\mathcal{T}(\mathbb{D}_C)$ are centered normal random variables with $\mathrm{var}\{\mathcal{T}(\mathbb{D}_C)\} \leq \mathrm{var}\{\mathcal{T}(\mathbb{D}_C^R)\}$. It follows that there exists $x_0 > 0$ such that, for any $x > x_0$,

$$\mathrm{P}\{|\mathcal{T}(\mathbb{D}_C)| > x\} \leq \mathrm{P}\{|\mathcal{T}(\mathbb{D}_C^R)| > x\}. \tag{30}$$

Now, assume that $H_{0,m} \cap H_{1,c}$ holds with $H_{1,c}$ defined in (20), $k^\star = \lfloor nt \rfloor$ for some $t \in (0,1)$ and that the copulas C_1 and C_2 are such that $\int_{[0,1]^2} \{C_1(\boldsymbol{u}) - C_2(\boldsymbol{u})\} \mathrm{d}\boldsymbol{u} \neq 0$. Then, from the proof of Proposition 3 of Bücher and Ruppert (2013) and from Proposition 5 (i) in this work, both $n^{-1/2}\mathbb{D}_n^R$ and $n^{-1/2}\mathbb{D}_n$ converge in probability in $\ell^\infty([0,1]^3)$ to the constant function K_t defined in Proposition 5 (i), and, by the continuous mapping theorem, both $n^{-1/2}\mathcal{T}(\mathbb{D}_n^R)$ and $n^{-1/2}\mathcal{T}(\mathbb{D}_n)$ converge in probability to $\mathcal{T}(K_t) \neq 0$. From the previous fact and (30), we can therefore expect that, as n increases, the asymptotic p-value for the test based on $\mathcal{T}(\mathbb{D}_n)$ will tend to be smaller than the asymptotic p-value for the test based on $\mathcal{T}(\mathbb{D}_n^R)$.

Unfortunately, at this point (see the discussion in Genest and Segers, 2010, Section 4), the previous argumentation cannot be extended to larger dimension or dependence conditions other than left-tail-decreasingness. Even for bivariate copulas that are LTD in both arguments, the case of the statistic S_n defined in (24) and its analogue S_n^R defined in (27) cannot be treated similarly, nor does it seem easy to deal with p-values computed approximately from multiplier replicates as in (25). Still, in our numerical experiments to be partially reported in the next section, the two versions of the test based on S_n considered in Section 3.1 were found to be consistently more powerful than the analogue multiplier test based on S_n^R.

Let us end this subsection with a remark. Assume that $H_{0,m}$ defined in (6) holds, and, for a moment, that we are in the unrealistic situation in which the continuous marginal c.d.f.s $F_1, \dots, F_d$ of the $\boldsymbol{X}_i$ are known. In this case, the sample $\boldsymbol{U}_1, \dots, \boldsymbol{U}_n$ with $\boldsymbol{U}_i = (F_1(X_{i1}), \dots, F_d(X_{id}))$ can be computed and it might seem intuitively better to use, instead of the process $\mathbb{D}_n$ defined in (21), the process $\tilde{\mathbb{D}}_n^\#$ defined as the process $\mathbb{D}_n^\#$ in (8) but with the $\boldsymbol{X}_i$ replaced by the $\boldsymbol{U}_i$. Using rather classical arguments, it can be verified that the difference between $\tilde{\mathbb{D}}_n^\#$ and $\mathbb{D}_n^R$ defined in (26) converge uniformly in probability to zero, which implies that, under H_0, the two processes converge weakly to the same limit $\mathbb{D}_C^R$ defined previously in this subsection. Corollary 4 then suggests that, at least in some scenarios, it is preferable to ignore the fact that the true marginal c.d.f.s are known and still use the process $\mathbb{D}_n$ for testing. Besides Proposition 1 of Genest and Segers (2010), yet another illustration of this phenomenon is given in Bücher (2012).

4 Monte Carlo experiments

4.1 The setting

Large-scale Monte Carlo experiments were carried out in order to study the finite-sample performance of various tests for change-point detection based on the empirical process $\mathbb{D}_n$ defined in (21). In the rest of the paper, the two versions of the test based on the statistic S_n defined in (24) will be referred to as *the test based on* $\hat{S}_n$ *(resp.* $\check{S}_n$*)* when approximate p-values are computed using the multiplier processes $\hat{\mathbb{D}}_n^{(m)}$ (resp. $\check{\mathbb{D}}_n^{(m)}$). As explained in the previous section, in addition to the tests based on the maximally selected Cramér–von Mises statistics $\hat{S}_n$ and $\check{S}_n$, we also considered tests based on Kolmogorov–Smirnov and Kuiper change-point statistics. Empirical results for these tests are not reported as they appear to be consistently less powerful than those based on $\hat{S}_n$ and $\check{S}_n$.

The two aforementioned tests were also compared with the test based on the statistic S_n^R defined in (27) derived from the process $\mathbb{D}_n^R$ defined in (26). An approximate p-value for the latter was computed using the multiplier processes

$$\mathbb{D}_n^{R,(m)}(s,\boldsymbol{u}) = \check{\mathbb{Z}}_n^{(m)}(s,\boldsymbol{u}) - \lambda_n(s)\check{\mathbb{Z}}_n^{(m)}(1,\boldsymbol{u}), \qquad (s,\boldsymbol{u}) \in [0,1]^{d+1},\ m \in \{1,\dots,M\},$$

where $\check{\mathbb{Z}}_n^{(m)}$ is defined in (16); see also Rémillard (2010, Section 5.2) and Bücher and Ruppert (2013, Section 3.2).

As a second competitor to the tests based on $\hat{S}_n$ and $\check{S}_n$, we included a test based on differences of sample Kendall's taus. Such a test was initially studied in Gombay and Horváth (1999, Section 3) in dimension two. Starting from the work of Gombay and Horváth (2002), multivariate extensions of this approach were recently investigated in Quessy et al. (2013). Given a d-dimensional random vector $\boldsymbol{X}$ with continuous marginal c.d.f.s $F_1,\dots,F_d$ and copula C, a natural multivariate extension of Kendall's tau is given by $\tau = (2^d \mathrm{E}\{C(\boldsymbol{U})\} - 1)/(2^{d-1}-1)$, where $\boldsymbol{U} = (F_1(X_1),\dots,F_d(X_d))$ (see e.g. Joe, 1997). One of the most powerful tests investigated in Quessy et al. (2013) is based on the statistic

$$\kappa_n = \max_{2\le k\le n-2} \frac{k(n-k)}{n^{3/2}} |\tau_k - \tau^\star_{n-k}|, \tag{31}$$

where τ_k and $\tau^\star_{n-k}$ are multivariate sample Kendall's taus computed from $\boldsymbol{X}_1,\dots,\boldsymbol{X}_k$ and $\boldsymbol{X}_{k+1},\dots,\boldsymbol{X}_n$, respectively. An approximate p-value for the latter test was obtained from multiplier replicates computed as explained in Gombay and Horváth (2002, page 251) with kernel h defined as $h(\boldsymbol{X}_i,\boldsymbol{X}_j) = \mathbf{1}(\boldsymbol{X}_i < \boldsymbol{X}_j) + \mathbf{1}(\boldsymbol{X}_i > \boldsymbol{X}_j)$ and up to the normalizing constant $2^d/(2^{d-1}-1)$.

To study the finite-sample performance of the tests based on $\hat{S}_n$, $\check{S}_n$, S_n^R and κ_n, data were first generated under H_0 defined in (5), then under $H_{0,m} \cap H_{1,c}$, where $H_{0,m}$ is defined in (6) and $H_{1,c}$ is defined in (20), and finally under $H_{1,m} \cap H_{0,c}$, where $H_{0,c}$ is defined in (7) and

$$\begin{aligned} H_{1,m} : {}& \exists \text{ distinct } F_{1,1}, F_{1,2} \text{ as well as } F_2,\dots,F_d \text{ and } k_1^\star \in \{1,\dots,n-1\} \\ & \text{such that } \boldsymbol{X}_1,\dots,\boldsymbol{X}_{k_1^\star} \text{ have marginal c.d.f.s } F_{1,1}, F_2,\dots,F_d \\ & \text{and } \boldsymbol{X}_{k_1^\star+1},\dots,\boldsymbol{X}_n \text{ have marginal c.d.f.s } F_{1,2}, F_2,\dots,F_d. \end{aligned} \tag{32}$$



Figure 1: The left (resp. right) plot shows 10,000 realizations of $aC_{\alpha,\beta}$ defined in (33) when C is the bivariate normal copula with parameter 0.97 and $(\alpha, \beta) = (0.25, 1)$ (resp. $(\alpha, \beta) = (0.5, 1)$).

In terms of copula families, we considered the Clayton, Gumbel–Hougaard, Frank and exchangeable normal families. A nice overview of their properties is given for instance in McNeil et al. (2005, Chapter 5). To fix parameter values for data generation, we exploited the well-known one-to-one relationship between Kendall's tau and the parameter value in the bivariate versions of these models. To increase the diversity of shapes in dimension two, asymmetric versions of the previous copulas were defined using the simplest form of Khoudraji's device (Khoudraji, 1995; Genest et al., 1998; Liebscher, 2008). Given an exchangeable bivariate copula C, Khoudraji's device can be used to define an asymmetric version of it as

$$aC_{\alpha,\beta}(u, v) = u^{1-\alpha}v^{1-\beta}C(u^{\alpha}, v^{\beta}), \qquad u, v \in [0, 1], \tag{33}$$

for arbitrary choices of $\alpha \neq \beta$, $\alpha, \beta \in (0, 1]$. As an illustration, the left (resp. right) plot of Figure 1 shows 10,000 realizations of $aC_{\alpha,\beta}$ when C is the bivariate normal copula with parameter 0.97 and $(\alpha, \beta) = (0.25, 1)$ (resp. $(\alpha, \beta) = (0.5, 1)$).

Two choices were considered for the multipliers: Rademacher random variables and standard normal random variables. The two settings led to very similar results suggesting that Theorem 1 (ii) might actually hold under Condition ($\mathcal{M}1$). The results reported in the forthcoming tables are those obtained with standard normal multipliers.

For each data generating scenario, the rejection rates of the tests were estimated from 1000 samples of size $n \in \{50, 100, 200\}$ and dimension $d = 2$ (and $d = 3$ when relevant) generated using the `copula` package (Hofert et al., 2012) for the R statistical system (R Development Core Team, 2012). All approximate p-values were computed from $M = 1000$ multiplier realizations and the tests were carried out at the 5% level of significance.

Table 1: Percentage of rejection of H_0 defined in (5) computed from 1000 random samples of size $n \in \{50, 100, 200\}$ generated under H_0, where C is either the d-dimensional Clayton (Cl), the Gumbel–Hougaard (GH) or the normal (N) copula whose bivariate margins have a Kendall's tau of τ.

			Cl				GH				N			
d	n	τ	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n
2	50	0.00	6.2	4.0	4.6	5.8	5.4	2.9	4.5	3.6	7.3	3.4	4.8	5.1
		0.25	6.7	6.2	5.6	5.6	5.5	3.3	5.4	4.3	4.4	3.0	6.3	3.4
		0.50	5.6	7.9	6.0	3.1	4.4	3.3	4.6	3.6	4.4	5.3	4.9	2.9
		0.75	6.0	16.6	5.5	3.2	3.2	6.7	4.3	2.6	3.6	9.1	4.9	2.9
	100	0.00	4.9	3.5	5.3	5.2	5.2	4.1	5.5	5.8	4.3	2.8	5.5	3.8
		0.25	6.1	6.6	5.0	5.0	5.0	3.3	6.2	5.6	5.3	4.0	5.5	5.3
		0.50	4.4	9.3	5.9	4.1	3.7	2.8	5.7	3.6	3.1	3.4	5.3	2.8
		0.75	2.7	10.0	4.6	2.1	2.5	4.7	4.4	1.8	2.1	6.0	5.6	1.5
	200	0.00	4.0	3.5	5.2	4.4	4.3	4.0	5.2	4.4	5.4	4.9	4.3	5.3
		0.25	4.7	5.2	6.3	4.9	3.3	3.0	3.8	4.6	4.0	3.9	5.2	5.2
		0.50	5.1	8.5	4.9	5.0	3.2	2.3	4.5	4.4	4.0	4.7	4.8	4.2
		0.75	2.6	9.3	5.9	3.2	1.5	3.1	5.2	3.1	1.9	4.8	5.7	1.5
3	50	0.00	4.3	1.5	3.0	2.6	4.2	2.1	3.6	3.4	5.5	2.8	3.4	4.4
		0.25	6.3	5.0	5.1	5.6	5.5	1.0	5.1	4.9	5.3	3.0	4.3	5.2
		0.50	7.2	8.1	5.9	6.2	2.7	0.9	5.7	4.1	3.0	2.2	4.6	3.6
		0.75	2.0	2.9	6.9	3.1	0.5	0.4	6.3	1.9	1.1	1.3	4.1	1.9
	100	0.00	4.5	3.4	4.5	4.3	4.5	2.8	4.6	3.7	4.5	2.7	3.9	5.0
		0.25	5.0	5.1	5.4	5.6	4.2	2.6	4.4	5.8	5.4	3.5	4.5	5.1
		0.50	5.7	7.6	6.3	5.0	3.3	1.3	5.0	4.6	3.2	3.1	3.9	3.6
		0.75	2.5	4.9	5.0	2.3	1.0	1.0	5.2	2.6	0.8	1.6	5.5	2.0
	200	0.00	3.3	2.5	4.3	4.1	3.5	3.2	4.3	4.1	4.8	4.0	4.7	5.2
		0.25	6.6	7.1	5.5	6.1	5.0	3.3	4.5	5.0	4.8	4.1	5.0	5.2
		0.50	6.0	9.2	4.5	5.6	3.0	2.4	5.9	5.3	4.8	4.3	4.8	5.5
		0.75	2.9	6.4	6.4	4.1	0.7	0.9	3.8	3.6	1.3	2.2	4.9	3.9

4.2 Rejection percentages under H_0

Table 1 reports the rejection percentages of H_0 defined in (5) computed from 1000 random samples generated under H_0, where the copula of F is either the Clayton, the Gumbel–Hougaard or the normal whose bivariate margins have a Kendall's tau of $\tau \in \{0, 0.25, 0.5, 0.75\}$. Since all the tests under consideration are rank-based, the marginal c.d.f.s were arbitrarily taken to be standard uniform. As one can see, the empirical levels of the tests based on $\hat{S}_n$ and κ_n are reasonably close to the 5% significance level, except sometimes for strongly dependent samples ($\tau = 0.75$) in which case the tests are, overall, too conservative. The test based on $\check{S}_n$ displays worryingly high empirical levels for samples generated from the Clayton copula when $\tau \geq 0.5$ (in particular for $d = 2$), although, as expected from the theoretical results of the previous section, the situation improves as n increases. It is on the contrary too conservative, overall, for

Table 2: Percentage of rejection of H_0 computed from 1000 samples of size $n \in \{50, 100, 200\}$ generated under $H_{0,m} \cap H_{1,c}$, where $H_{1,c}$ is defined in (20), $k^\star = \lfloor nt \rfloor$, C_1 and C_2 are both either bivariate Clayton (Cl), Gumbel-Hougaard (GH) or normal (N) copulas such that C_1 has a Kendall's tau of 0.2 and C_2 a Kendall's tau of τ.

			Cl				GH				N			
n	τ	t	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n
50	0.4	0.10	7.5	8.1	5.7	5.8	6.1	3.7	4.3	5.5	6.1	4.8	4.3	4.6
		0.25	12.1	10.6	4.0	8.4	9.4	5.1	5.0	8.8	10.4	7.6	5.5	7.7
		0.50	18.0	16.1	6.3	13.3	12.1	7.8	4.8	12.5	12.3	8.4	5.8	10.5
	0.6	0.10	14.5	17.0	5.4	11.2	11.4	7.7	6.2	9.9	11.4	9.8	7.2	10.2
		0.25	35.5	34.4	7.4	28.0	29.9	21.4	6.4	27.5	31.3	21.4	7.1	29.2
		0.50	47.3	41.6	7.0	43.7	45.3	30.3	8.9	46.1	46.0	33.9	9.1	45.1
100	0.4	0.10	7.1	8.7	6.1	6.7	6.6	5.1	5.1	7.0	5.8	5.2	5.3	6.0
		0.25	18.8	19.9	5.2	16.1	16.9	13.2	5.8	18.7	14.9	12.5	6.1	16.9
		0.50	26.5	26.4	7.3	26.7	23.8	18.8	7.3	27.4	22.6	19.1	7.9	24.8
	0.6	0.10	21.5	25.1	5.8	19.6	16.7	12.0	5.1	18.3	17.5	16.9	6.1	19.1
		0.25	65.1	66.0	6.3	63.1	61.2	51.5	7.5	64.7	62.9	54.8	9.9	67.4
		0.50	82.1	81.6	14.7	83.8	78.8	69.7	14.9	83.3	79.1	73.7	11.8	86.0
200	0.4	0.10	11.1	13.8	5.8	11.1	8.3	8.1	5.5	9.2	9.5	9.8	5.0	11.7
		0.25	30.8	33.9	5.9	31.5	27.6	24.8	6.4	31.6	29.6	28.3	6.8	34.8
		0.50	47.1	48.6	9.0	51.3	45.8	41.4	8.7	50.4	47.1	46.1	9.4	54.0
	0.6	0.10	36.4	41.3	6.8	40.7	34.4	31.7	7.1	44.0	36.0	36.3	6.7	49.3
		0.25	92.6	93.2	12.3	93.8	91.4	88.9	16.7	95.0	91.3	90.2	12.0	95.5
		0.50	98.9	99.3	22.2	99.5	98.5	98.1	22.0	99.6	99.3	99.1	21.1	100.0

samples generated from the Gumbel–Hougaard copula. The test based on S_n^R appears to hold its level reasonably well in all the scenarios under consideration.

To investigate the influence of negative dependence, a similar setting was considered in dimension two for normal and Frank copulas whose parameters were fixed so that the copulas have a Kendall's tau of $\tau \in \{-0.25, -0.5, -0.75\}$. The results (not reported) are similar to those obtained in the case of positive dependence. Finally, the empirical levels of the tests were computed from bivariate datasets of size $n \in \{50, 100, 200\}$ generated from asymmetric copulas obtained as in (33) when C is the bivariate normal copula with parameter 0.97, $\alpha \in \{0.25, 0.5\}$ and $\beta = 1$ (see also Figure 1). The resulting rejection percentages (not reported) appear to be reasonably close to the 5% significance level in all scenarios.

4.3 Rejection percentages under $H_{0,m} \cap H_{1,c}$

The aim of the first experiment under $H_{0,m} \cap H_{1,c}$ was to investigate the influence of a change in the strength of the dependence only on the power of the tests. Table 2 reports the rejection percentages of H_0 computed from 1000 random samples generated

Table 3: Percentage of rejection of H_0 computed from 1000 samples of size $n \in \{50, 100, 200\}$ generated under $H_{0,m} \cap H_{1,c}$, where $H_{1,c}$ is defined in (20), $k^\star = \lfloor nt \rfloor$, C_1 and C_2 are both either bivariate normal (N) or Frank (F) copulas such that C_1 has a Kendall's tau of -0.2 and C_2 a Kendall's tau of τ.

			N				F			
n	τ	t	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n
50	-0.4	0.10	5.1	3.3	3.7	4.2	6.1	4.0	4.1	4.8
		0.25	9.5	6.2	4.2	7.8	12.8	8.8	5.1	9.2
		0.50	15.1	10.0	6.6	12.1	14.7	11.1	6.4	11.2
	-0.6	0.10	10.9	8.6	3.7	8.9	10.3	7.2	4.0	8.4
		0.25	33.6	23.8	6.5	26.7	36.2	26.4	5.1	29.3
		0.50	48.6	36.5	6.9	47.6	50.3	37.6	8.4	47.5
100	-0.4	0.10	6.6	5.3	4.2	6.5	6.9	6.0	4.0	6.3
		0.25	18.4	15.9	5.8	18.5	17.3	15.5	6.1	17.8
		0.50	24.2	21.3	7.0	25.1	28.6	25.5	7.9	30.7
	-0.6	0.10	15.9	15.7	6.1	18.9	18.5	15.6	4.9	21.2
		0.25	59.9	54.3	11.6	65.6	66.7	59.8	12.6	69.1
		0.50	81.9	76.7	19.2	89.0	84.9	80.9	20.6	87.8
200	-0.4	0.10	6.9	7.4	5.3	9.2	10.1	9.5	5.2	11.1
		0.25	29.1	27.8	7.4	33.6	29.4	28.2	8.3	33.0
		0.50	46.4	44.9	11.0	52.5	50.6	48.2	13.0	55.7
	-0.6	0.10	37.3	37.7	10.0	48.6	40.6	40.5	9.8	52.9
		0.25	93.8	92.8	30.0	96.9	96.0	94.6	30.1	97.7
		0.50	99.4	99.2	38.9	99.7	99.6	99.5	48.5	99.9

under $H_{0,m} \cap H_{1,c}$, where $H_{0,m}$ is defined in (6) and $H_{1,c}$ is defined in (20), $k^\star = \lfloor nt \rfloor$ for $t \in \{0.1, 0.25, 0.5\}$, and C_1 and C_2 are both either bivariate Clayton, Gumbel–Hougaard or normal copulas such that C_1 has a Kendall's tau of 0.2 and C_2 a Kendall's tau of $\tau \in \{0.4, 0.6\}$. As one can see, the tests based on $\hat{S}_n$, $\check{S}_n$ and κ_n consistently outperform the test based on S_n^R whose power is relatively weak for the sample sizes under consideration. The test based on $\hat{S}_n$ appears overall slightly more powerful than its two competitors for $n = 50$, while, for $n = 200$, the highest rejection rates are obtained, overall, for the test based on κ_n. Unsurprisingly, the estimated powers are the highest when the change occurs in the middle of the sequence. The conclusions remain very similar in dimension three (results not reported) except that all the rejection rates are higher. This could have been expected as an increase in dimension can be regarded as an increase in sample size for data generated from exchangeable copulas.

As the theoretical hints of Section 3.2 were provided for copulas with positive dependence, it seemed important to empirically investigate the case of negative dependence. Table 3 is the analogue of Table 2 in that situation. This time C_1 and C_2 are both either bivariate normal or Frank copulas such that C_1 has a Kendall's tau of -0.2 and C_2 a Kendall's tau of $\tau \in \{-0.4, -0.6\}$. While the rejection percentages of the tests based on $\hat{S}_n$, $\check{S}_n$ and κ_n remain comparable with those of Table 2, we see that, somehow

Table 4: Percentage of rejection of H_0 computed from 1000 samples of size $n \in \{100, 200\}$ generated under $H_{0,m} \cap H_{1,c}$, where $H_{1,c}$ is defined in (20), $k^\star = \lfloor nt \rfloor$, C_1 (resp. C_2) is a d-dimesional Clayton (resp. Gumbel-Hougaard) copula whose bivariate margins have a Kendall's tau of τ.

			$d = 2$				$d = 3$			
n	τ	t	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n
100	0.25	0.25	5.7	4.1	5.3	6.5	5.3	3.1	4.4	5.0
		0.50	6.2	5.7	5.6	4.6	9.1	6.0	5.8	7.7
		0.75	6.6	6.3	3.5	5.9	5.8	5.4	4.9	4.8
	0.50	0.25	5.5	5.9	5.8	3.2	4.6	2.9	5.0	4.3
		0.50	10.5	12.2	5.1	4.5	15.1	15.1	6.8	3.3
		0.75	8.3	11.9	4.4	4.5	7.7	9.9	5.3	4.1
	0.75	0.25	4.0	7.6	5.1	2.0	2.5	1.9	4.3	2.0
		0.50	12.5	19.9	6.0	2.0	9.8	13.2	5.2	2.3
		0.75	8.2	16.4	6.5	1.0	4.8	6.7	3.8	1.9
200	0.25	0.25	5.8	5.3	6.1	6.9	5.8	3.7	5.8	5.7
		0.50	8.5	8.7	5.4	5.2	9.8	9.7	6.6	6.2
		0.75	6.4	6.9	5.5	5.2	9.0	9.1	5.3	6.4
	0.50	0.25	10.4	11.5	6.8	3.7	12.6	10.1	6.7	3.7
		0.50	30.9	37.5	5.3	5.6	44.0	45.8	7.1	6.0
		0.75	16.3	23.2	6.1	4.1	20.1	27.0	5.1	4.5
	0.75	0.25	11.3	18.0	4.9	2.5	15.6	16.7	7.3	2.5
		0.50	43.4	54.4	6.2	2.5	58.9	63.1	5.2	2.5
		0.75	21.3	36.4	4.6	2.5	23.6	36.0	6.8	4.7

surprisingly, those of the test based on S_n^R are higher, though still substantially smaller than those of the three other tests.

The next experiment under $H_{0,m} \cap H_{1,c}$ consisted of taking C_1 and C_2 from different copula families but such that their bivariate margins have the same value $\tau \in \{0.25, 0.5, 0.75\}$ of Kendall's tau. In other words, the aim was to investigate the influence of a change in the shape of the dependence on the power of the tests. Because the test based on κ_n only measures a difference in dependence strength, it has no power against this type of alternatives. Note that this is a hard problem, as distinguishing copulas on the basis of low amounts of data is known to be difficult, for instance, from the corresponding goodness-of-fit testing literature (see e.g. Genest et al., 2009; Kojadinovic and Yan, 2011). The fact that the change-point is unknown makes the problem under consideration even more difficult. Two of the most easily distinguishable copulas are the Clayton and the Gumbel–Hougaard because the former has lower tail dependence while the latter has upper tail dependence (see e.g. McNeil et al., 2005, Chapter 5). The rejection percentages for this scenario are reported in Table 4. As one can see, for the sample sizes under consideration, the test based on S_n^R has hardly any power against this type of alternative. For the tests based on $\hat{S}_n$ and $\check{S}_n$, the highest rejection percentages are obtained for $\tau = 0.75$. Overall, the tests based on $\hat{S}_n$ and $\check{S}_n$ appear to have roughly

Table 5: Percentage of rejection of H_0 computed from 1000 samples of size $n \in \{50, 100, 200\}$ generated under $H_{0,m} \cap H_{1,c}$, where $H_{1,c}$ is defined in (20), $k^\star = \lfloor nt \rfloor$, and C_1 and C_2 are asymmetric copulas defined as in (33) from the normal copula with parameter 0.97. For C_1, $(\alpha, \beta) = (1, \delta)$ and for C_2, $(\alpha, \beta) = (\delta, 1)$, with $\delta \in \{0.25, 0.5\}$.

n	t	δ	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n
100	0.25	0.25	6.0	4.3	5.9	4.1
		0.50	7.9	7.1	4.4	5.4
	0.50	0.25	10.3	8.4	4.4	5.2
		0.50	11.4	10.3	6.2	4.6
200	0.25	0.25	10.5	9.6	5.4	5.6
		0.50	14.1	16.3	5.7	6.3
	0.50	0.25	23.8	23.8	6.9	5.3
		0.50	50.7	59.4	8.1	4.8

equivalent power for $\tau = 0.25$. The fact that the rejection rates of the test based on $\check{S}_n$ are sometimes higher than those of the test based on $\hat{S}_n$ when $\tau \geq 0.5$ should be interpreted with care as the former was found to be too liberal in the case of strongly dependent data in the previous subsection. Notice also that the results are not symmetric with respect to $t = 0.5$. Indeed, from the results for $t = 0.25$ and $t = 0.75$, we see that it seems easier to reject H_0 when the largest of the two subsamples comes from the Clayton copula. Finally, we see that the estimated rejection rates tend overall to increase as d changes from 2 to 3, which could be explained, as earlier, from the exchangeability of the copulas.

Similar experiments were carried for C_1 and C_2 in other families. When one of the copulas was taken from the Clayton family, the tests based on $\hat{S}_n$ and $\check{S}_n$ displayed some power for $\tau = 0.75$. In other situations, such as when C_1 is from the normal family and C_2 is from the Frank family, the tests had hardly any power for the sample sizes under consideration. Again, this is not surprising from empirical results obtained in the context of goodness-of-fit testing.

As a last experiment under $H_{0,m} \cap H_{1,c}$, we considered the situation in which C_1 and C_2 are asymmetric copulas defined as in (33) from the normal copula with parameter 0.97. For C_1, we took $(\alpha, \beta) = (1, \delta)$ and for C_2, $(\alpha, \beta) = (\delta, 1)$, with $\delta \in \{0.25, 0.5\}$, so that C_1 and C_2 have the same value of Kendall's tau but "opposite" asymmetry. The left (resp. right) plot of Figure 1 shows 10,000 realizations of C_2 for $\delta = 0.25$ (resp. $\delta = 0.5$). The results of the experiment are reported in Table 5. As expected, the power of the tests based on $\hat{S}_n$ and $\check{S}_n$ are the largest when the asymmetry is the strongest ($\delta = 0.5$). Again, for the sample sizes under consideration, the test based on S_n^R has hardly any power against this type of alternative.

4.4 Rejection percentages under $H_{0,c} \cap H_{1,m}$

We concluded our Monte Carlo experiments by an investigation of the power of the tests for distributional changes solely due to a break in the first marginal c.d.f. Samples of size

$n \in \{50, 100, 200\}$ were generated under $H_{0,c} \cap H_{1,m}$, where $H_{1,m}$ is defined in (32), such that the $k_1^\star = \lfloor nt_1 \rfloor$ first observations of each sample were from a d-dimensional c.d.f. with copula C and $N(0,1)$ margins, and the $n - \lfloor nt_1 \rfloor$ last observations were from a d-dimensional c.d.f. with copula C whose first margin is $N(\mu, 1)$ and whose d-1 remaining margins are $N(0,1)$. The values 0.25 and 0.5 (resp. 0.5, 1 and 2) were considered for t_1 (resp. μ). The rejection percentages of H_0 when C is a normal copula are partially reported in Table 6. Similar results were obtained for the other copula families considered in the simulations. As one can see, the test based on S_n^R is this time substantially more powerful than those based on $\hat{S}_n$, $\check{S}_n$ and κ_n. This was expected in Section 3.2 because of (28), which shows that the test based on S_n^R is actually a general-purpose test for detecting distributional changes. As far as the tests based on $\hat{S}_n$, $\check{S}_n$ and κ_n are concerned, their overall weak power against this type of alternatives can be explained by the fact that they were designed for detecting changes in the copula. Their rejection rates are close to the 5% significance level for $\mu = 0.5$ and are the highest for $(\mu, t_1) = (2, 0.25)$ and $\tau = 0.5$. Notice that another reading of the results for $\mu = 0.5$ is that the tests based on $\hat{S}_n$, $\check{S}_n$ and κ_n, regarded as procedures for testing directly $H_{0,c}$, seem relatively robust against small breaks in mean in one margin.

5 Concluding remarks

The weak limit of the sequential empirical copula process $\mathbb{C}_n$ defined in (3) was obtained and the asymptotic validity of a multiplier bootstrap was established under the nonrestrictive smoothness conditions considered by Segers (2012). These theoretical results were then used to derive tests for change-point detection particularly sensitive to changes in the copula. Among several tests derived from the process $\mathbb{D}_n$ defined in (21), the one based on the maximally selected Cramér–von Mises statistic S_n defined in (24) was found to be the most powerful in finite-samples. It was compared with the analogue test based on the statistic S_n^R defined in (27) derived from the process $\mathbb{D}_n^R$ defined in (26) considered in Rémillard (2010, Section 5.2) and Bücher and Ruppert (2013, Section 3.2). Theoretical hints suggesting that, at least in some scenarios, tests based on $\mathbb{D}_n$ should be more powerful for detecting changes in the copula than tests based on $\mathbb{D}_n^R$ were provided, and this was confirmed by extensive Monte Carlo experiments. In these experiments, the test based on $\hat{S}_n$ and the one based on κ_n defined in (31) displayed, overall, the best finite-sample behavior. The latter was found, overall, to be the most powerful for detecting changes in the dependence strength. The former has however the advantage of being consistent against all alternatives under $H_{0,m} \cap H_{1,c}$, where $H_{0,m}$ is defined in (6) and $H_{1,c}$ is defined in (20).

It is our understanding that none of the existing nonparametric change-point tests designed to be particularly sensitive to changes in the copula can be used to directly test $H_{0,c}$ defined in (7) unless $H_{0,m}$ defined in (6) is assumed to hold. If such tests are applied and H_0 defined in (5) is rejected, a pragmatic approach would then be to additionally perform d univariate change-point tests on each of the component sequences. Should there be no evidence against $H_{0,m}$, rejection of $H_{0,c}$ could be concluded.

The latter approach is however no completely satisfactory and a future research direction would be to design nonparametric change-point tests that can be used to test $H_{0,c}$ whether $H_{0,m}$ holds or not (see also the discussion in Quessy et al., 2013, Section 4). Another important future contribution would be to extend the approach considered in this work to the case of serially dependent observations. For that purpose, a good starting point is the work of Bücher and Ruppert (2013).

Acknowledgments

The authors are very grateful to Axel Bücher, Mark Holmes, Jean-François Quessy, Martin Ruppert and Johan Segers for fruitful discussions.

A Proofs of Propositions 1 and 2

Proof of Proposition 1. To show the desired result, let us first show that

$$A_n = \sup_{s \in [0,1]} \sup_{\boldsymbol{u} \in [0,1]^d} \left| \mathbb{C}_n(s, \boldsymbol{u}) - \left\{ \mathbb{Z}_n(s, \boldsymbol{u}) - \sum_{j=1}^{d} C^{[j]}(\boldsymbol{u}) \mathbb{Z}_n(s, \boldsymbol{u}^{(j)}) \right\} \right| \xrightarrow{\text{a.s.}} 0. \tag{34}$$

It can be verified that A_n can be rewritten as $A_n = \max_{1 \leq k \leq n} \sqrt{k/n} B_k$, where

$$B_k = \sup_{\boldsymbol{u} \in [0,1]^d} \left| \mathbb{C}_k(1, \boldsymbol{u}) - \left\{ \mathbb{Z}_k(1, \boldsymbol{u}) - \sum_{j=1}^{d} C^{[j]}(\boldsymbol{u}) \mathbb{Z}_k(1, \boldsymbol{u}^{(j)}) \right\} \right|. \tag{35}$$

We can further write $A_n = \max\{\sqrt{1/n} B_1, \sqrt{2/n} B_2, \max_{3 \leq k \leq n} \sqrt{k/n} B_k\}$. To show that $A_n \xrightarrow{\text{a.s.}} 0$, it therefore suffices to show that $\max_{3 \leq k \leq n} \sqrt{k/n} B_k \xrightarrow{\text{a.s.}} 0$.

Let $a_n = n^{-1/4} (\log n)^{1/2} (\log \log n)^{1/4}$, $n \geq 3$. Then,

$$\max_{3 \leq k \leq n} \sqrt{\frac{k}{n}} B_k \leq \max_{3 \leq k \leq n} \sqrt{\frac{k}{n}} a_k \times \max_{3 \leq k \leq n} \frac{B_k}{a_k}.$$

First, notice that

$$\max_{3 \leq k \leq n} \sqrt{\frac{k}{n}} a_k = n^{-1/2} \max_{3 \leq k \leq n} k^{1/4} (\log k)^{1/2} (\log \log k)^{1/4} = a_n \to 0.$$

Next, (12), which is equivalent to $\limsup B_n / a_n < \infty$ with probability one, implies that $\max_{3 \leq k \leq n} B_k / a_k \leq \sup_{k \geq 3} B_k / a_k < \infty$ almost surely, from which we obtain that $\max_{3 \leq k \leq n} \sqrt{k/n} B_k \xrightarrow{\text{a.s.}} 0$. The desired result is finally a consequence of the weak convergence of $\mathbb{Z}_n$ to $\mathbb{Z}_C$ in $\ell^\infty([0,1]^{d+1})$ and the continuous mapping theorem. ■

Proof of Proposition 2. The proof starts as the proof of Lemma 2 of Kojadinovic et al. (2011). Without loss of generality, we deal with $C_n^{[1]}$. For any $\boldsymbol{u} \in [0,1]^d$, let $\boldsymbol{u}_{-1}$ denote the vector $(u_2, \ldots, u_d)$ of $[0,1]^{d-1}$. Also, let $\delta > 0$ be a real number such that

$0 < \delta < a < b < 1-\delta < 1$, and let n be sufficiently large such that, for any $x \in [a,b]$, $x \pm n^{-1/2} \in [\delta, 1-\delta]$. It follows that, for any $u_1 \in [a,b]$, $u_{1,n}^+ = u_1 + n^{-1/2}$, $u_{1,n}^- = u_1 - n^{-1/2}$, and $u_{1,n}^+ - u_{1,n}^- = 2n^{-1/2}$. Also, for any $\boldsymbol{u} \in [0,1]^d$ such that $u_1 \in [a,b]$, we have that

$$\begin{aligned} \sup_{\substack{\boldsymbol{u}\in[0,1]^d \\ u_1\in[a,b]}} |C_n^{[1]}(\boldsymbol{u}) - C^{[1]}(\boldsymbol{u})| &\leq \sup_{\substack{\boldsymbol{u}\in[0,1]^d \\ u_1\in[a,b]}} \left| \frac{1}{2n^{-1/2}} \left\{ C(u_{1,n}^+, \boldsymbol{u}_{-1}) - C(u_{1,n}^-, \boldsymbol{u}_{-1}) \right\} - C^{[1]}(\boldsymbol{u}) \right| \\ &\quad + \sup_{\substack{\boldsymbol{u}\in[0,1]^d \\ u_1\in[a,b]}} \left| \mathbb{C}_n(1, u_{1,n}^+, \boldsymbol{u}_{-1}) - \mathbb{C}_n(1, u_{1,n}^-, \boldsymbol{u}_{-1}) \right|, \quad (36) \end{aligned}$$

where $\mathbb{C}_n(1,\cdot) = \sqrt{n}(C_n - C)$; see also (3). Since $C^{[1,1]}$ exists and is continuous on $V_1 = \{\boldsymbol{u} \in [0,1]^d : 0 < u_1 < 1\}$ according to Condition ($\mathcal{C}2$), for any $\boldsymbol{u} \in [a,b] \times [0,1]^{d-1}$, there exists $u_{1,n}^{+,*} \in (u_1, u_{1,n}^+) \subset [\delta, 1-\delta]$ and $u_{1,n}^{-,*} \in (u_{1,n}^-, u_1) \subset [\delta, 1-\delta]$, such that

$$C(u_{1,n}^+, \boldsymbol{u}_{-1}) = C(\boldsymbol{u}) + C^{[1]}(\boldsymbol{u})n^{-1/2} + C^{[1,1]}(u_{1,n}^{+,*}, \boldsymbol{u}_{-1})(2n)^{-1}$$

and

$$C(u_{1,n}^-, \boldsymbol{u}_{-1}) = C(\boldsymbol{u}) - C^{[1]}(\boldsymbol{u})n^{-1/2} + C^{[1,1]}(u_{1,n}^{-,*}, \boldsymbol{u}_{-1})(2n)^{-1},$$

which implies that

$$\sup_{\substack{\boldsymbol{u}\in[0,1]^d \\ u_1\in[a,b]}} \left| \frac{1}{2n^{-1/2}} \left\{ C(u_{1,n}^+, \boldsymbol{u}_{-1}) - C(u_{1,n}^-, \boldsymbol{u}_{-1}) \right\} - C^{[1]}(\boldsymbol{u}) \right| \leq n^{-1/2} \sup_{\boldsymbol{u}\in V_1^\delta} \left| C^{[1,1]}(\boldsymbol{u}) \right| = O(n^{-1/2}),$$

since $C^{[1,1]}$ is continuous on $V_1^\delta = \{\boldsymbol{u} \in V_1 : u_1 \in [\delta, 1-\delta]\} \subset V_1$.

For the second supremum on the right of (36), we start from (12) and write

$$\mathbb{C}_n(1, \boldsymbol{u}) = \mathbb{Z}_n(1, \boldsymbol{u}) - \sum_{j=1}^d C^{[j]}(\boldsymbol{u}) \mathbb{Z}_n(1, \boldsymbol{u}^{(j)}) + R_n(\boldsymbol{u}),$$

where $\sup_{\boldsymbol{u}\in[0,1]^d} |R_n(\boldsymbol{u})| = O\left(n^{-1/4}(\log n)^{1/2}(\log\log n)^{1/4}\right)$ almost surely. Then,

$$\begin{aligned} &\sup_{\substack{\boldsymbol{u}\in[0,1]^d \\ u_1\in[a,b]}} \left| \mathbb{C}_n(1, u_{1,n}^+, \boldsymbol{u}_{-1}) - \mathbb{C}_n(1, u_{1,n}^-, \boldsymbol{u}_{-1}) \right| \\ &\qquad \leq \sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in V_1^\delta, \boldsymbol{u}_{-1}=\boldsymbol{v}_{-1} \\ |u_1 - v_1| \leq 2n^{-1/2}}} |\mathbb{C}_n(1, \boldsymbol{u}) - \mathbb{C}_n(1, \boldsymbol{v})| \leq A_{1,n} + A_{2,n} + A_{3,n} + A_{4,n}, \end{aligned}$$

where

$$\begin{aligned} A_{1,n} &= \sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in V_1^\delta, \boldsymbol{u}_{-1}=\boldsymbol{v}_{-1} \\ |u_1 - v_1| \leq 2n^{-1/2}}} \left| \mathbb{Z}_n(1, \boldsymbol{u}) - \mathbb{Z}_n(1, \boldsymbol{v}) \right|, \\ A_{2,n} &= \sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in V_1^\delta, \boldsymbol{u}_{-1}=\boldsymbol{v}_{-1} \\ |u_1 - v_1| \leq 2n^{-1/2}}} \left| C^{[1]}(\boldsymbol{u}) \mathbb{Z}_n(1, \boldsymbol{u}^{(1)}) - C^{[1]}(\boldsymbol{v}) \mathbb{Z}_n(1, \boldsymbol{v}^{(1)}) \right|, \\ A_{3,n} &= \sum_{j=2}^d \sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in V_1^\delta, \boldsymbol{u}_{-1}=\boldsymbol{v}_{-1} \\ |u_1 - v_1| \leq 2n^{-1/2}}} \left| \left\{ C^{[j]}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{v}) \right\} \mathbb{Z}_n(1, \boldsymbol{u}^{(j)}) \right|, \end{aligned}$$

and $A_{4,n} = 2\sup_{\boldsymbol{u}\in[0,1]^d}|R_n(\boldsymbol{u})| = O\left(n^{-1/4}(\log n)^{1/2}(\log\log n)^{1/4}\right)$ almost surely. Proceeding as in the proof of Proposition 4.2 of Segers (2012) for the term I_n, it can be verified that, with probability one,

$$A_{1,n} \leq \sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in[0,1]^d \\ \forall j\, |u_j - v_j| \leq 2n^{-1/2}}} |\mathbb{Z}_n(1,\boldsymbol{u}) - \mathbb{Z}_n(1,\boldsymbol{v})| = O\left(n^{-1/4}(\log n)^{1/2}\right).$$

For $A_{2,n}$, we have $A_{2,n} \leq B_{1,n} + B_{2,n}$, where

$$B_{1,n} = \sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in V_1^\delta, \boldsymbol{u}_{-1}=\boldsymbol{v}_{-1} \\ |u_1 - v_1| \leq 2n^{-1/2}}} \left|C^{[1]}(\boldsymbol{u}) - C^{[1]}(\boldsymbol{v})\right| \times \sup_{\boldsymbol{u}\in V_1^\delta}\left|\mathbb{Z}_n(1,\boldsymbol{u}^{(1)})\right|$$

and

$$B_{2,n} = \sup_{\boldsymbol{u}\in V_1^\delta}\left|C^{[1]}(\boldsymbol{u})\right| \times A_{1,n} = O(n^{-1/4}(\log n)^{-1/2}) \qquad \text{almost surely.}$$

For $B_{1,n}$, from the mean value theorem, we obtain that, with probability one,

$$B_{1,n} \leq \sup_{\boldsymbol{u}\in V_1^\delta}\left|C^{[1,1]}(\boldsymbol{u})\right| \times \frac{\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_n(1,\boldsymbol{u}^{(1)})\right|}{(\log\log n)^{1/2}} \times 2n^{-1/2}(\log\log n)^{1/2} = O(n^{-1/2}(\log\log n)^{1/2}),$$

since $\sup_{n\geq 3}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_n(1,\boldsymbol{u}^{(1)})\right|/(\log\log n)^{1/2} < \infty$ almost surely by the law of the iterated logarithm for empirical c.d.f.s (see e.g. Chung, 1949, Theorem 2*). For the term $A_{3,n}$, we proceed as in the proof of Proposition 4.2 of Segers (2012) for the term III_n. Write $A_{3,n} = \sum_{j=2}^d A_{3,n,j}$, where

$$A_{3,n,j} = \sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in V_1^\delta, \boldsymbol{u}_{-1}=\boldsymbol{v}_{-1} \\ |u_1 - v_1| \leq 2n^{-1/2}}} \left|\left\{C^{[j]}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{v})\right\}\mathbb{Z}_n(1,\boldsymbol{u}^{(j)})\right|.$$

Let $\gamma_n = n^{-1/2}(\log n)(\log\log n)^{-1/2}$, $n \geq 3$, let $V_j^{\gamma_n} = \{\boldsymbol{u} \in V_j : u_j \in [\gamma_n, 1-\gamma_n]\} \subset V_j$, and let $\bar{V}_j^{\gamma_n} = [0,1]^d \setminus V_j^{\gamma_n}$. Then, using the fact that V_1^δ is the disjoint union of $V_1^\delta \cap V_j^{\gamma_n}$ and $V_1^\delta \cap \bar{V}_j^{\gamma_n}$, and that $\sup_{\boldsymbol{u},\boldsymbol{v}\in[0,1]^d}\left|C^{[j]}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{v})\right| \leq 1$, we obtain that $A_{3,n,j} \leq A'_{3,n,j} + A''_{3,n,j}$, where $A'_{3,n,j} = \sup_{\boldsymbol{u}\in\bar{V}_j^{\gamma_n}}\left|\mathbb{Z}_n(1,\boldsymbol{u}^{(j)})\right|$ and

$$A''_{3,n,j} = \sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in V_1^\delta\cap V_j^{\gamma_n}, \boldsymbol{u}_{-1}=\boldsymbol{v}_{-1} \\ |u_1 - v_1| \leq 2n^{-1/2}}} \left|\left\{C^{[j]}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{v})\right\}\mathbb{Z}_n(1,\boldsymbol{u}^{(j)})\right|.$$

As verified in Segers (2012, Eq. (4.3)) using Theorem 2 (iii) of Einmahl and Mason (1988), $A'_{3,n,j} = O\left(n^{-1/4}(\log n)^{1/2}(\log\log n)^{1/4}\right)$ almost surely. Also, from Segers (2012, Lemma 4.3), there exists a constant $\alpha > 0$ such that

$$A''_{3,n,j} \leq \alpha \sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in V_1^\delta\cap V_j^{\gamma_n}, \boldsymbol{u}_{-1}=\boldsymbol{v}_{-1} \\ |u_1 - v_1| \leq 2n^{-1/2}}} \left\{\frac{|u_1 - v_1|}{u_j(1-u_j)}\left|\mathbb{Z}_n(1,\boldsymbol{u}^{(j)})\right|\right\}.$$

Using the fact that $\{u_j(1-u_j)\}^{-1/2} \leq \{\gamma_n(1-\gamma_n)\}^{-1/2} = O(\gamma_n^{-1/2})$ for all $u_j \in [\gamma_n, 1-\gamma_n]$, and that, as verified in Segers (2012, page 775) for instance from Mason (1981, Theorem

2), $\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_n(1,\boldsymbol{u}^{(j)})\right|\{u_j(1-u_j)\}^{-1/2} = O\left((\log n)^{1/2}\log\log n\right)$ almost surely, we have, with probability one,

$$A''_{3,n,j} \leq \alpha 2n^{-1/2} \times \sup_{\boldsymbol{u}\in[0,1]^d} \frac{\left|\mathbb{Z}_n(1,\boldsymbol{u}^{(j)})\right|}{\{u_j(1-u_j)\}^{1/2}} \times \{\gamma_n(1-\gamma_n)\}^{-1/2} = O\left(n^{-1/4}(\log\log n)^{5/4}\right).$$

It follows that $A_{3,n} = O\left(n^{-1/4}(\log n)^{1/2}(\log\log n)^{1/4}\right)$ almost surely, which implies the desired result. ■

B Proof of Theorem 1

Let us first define additional notation. Let G_n be the empirical c.d.f. computed from the unobservable random sample $\boldsymbol{U}_1,\dots,\boldsymbol{U}_n$ and let $G_{n,j}$, $j\in\{1,\dots,d\}$, be the univariate margins of G_n. Recall that, for any $j\in\{1,\dots,d\}$, $R_{ij,n}$ is the rank of X_{ij} among $X_{1j},\dots,X_{nj}$, or, equivalently, the rank of U_{ij} among $U_{1j},\dots,U_{nj}$. We also have that, for any $i\in\{1,\dots,n\}$ and $j\in\{1,\dots,d\}$, $G_{n,j}(U_{ij}) = R_{ij,n}/n = \hat{U}_{ij,n}(n+1)/n$. Furthermore, for any $j\in\{1,\dots,d\}$, let

$$G_{n,j}^{-1}(u) = \inf\{v\in[0,1] : G_{n,j}(v)\geq u\}, \qquad u\in[0,1],$$

be the quantile function associated with $G_{n,j}$. It is well-known that

$$G_{n,j}^{-1}(u) = \begin{cases} U_{k:n,j} & \text{if } (k-1)/n < u \leq k/n, \\ 0 & \text{if } u = 0, \end{cases} \tag{37}$$

where $U_{1:n,j} < \dots < U_{n:n,j}$ are the order statistics associated with the jth coordinate sample $U_{1j},\dots,U_{nj}$. Next, we define

$$G_n^{-1}(\boldsymbol{u}) = \left(G_{n,1}^{-1}(u),\dots,G_{n,d}^{-1}(u)\right), \qquad \boldsymbol{u}\in[0,1]^d. \tag{38}$$

Finally, for any $m\in\{1,\dots,M\}$ and $(s,\boldsymbol{u})\in[0,1]^{d+1}$, let

$$\tilde{\tilde{\mathbb{Z}}}_n^{(m)}(s,\boldsymbol{u}) = \frac{1}{\sqrt{n}}\sum_{i=1}^{\lfloor ns\rfloor}\xi_i^{(m)}\left[\mathbf{1}\left\{\boldsymbol{U}_i\leq G_{\lfloor ns\rfloor}^{-1}(\boldsymbol{u})\right\} - G_{\lfloor ns\rfloor}\left\{G_{\lfloor ns\rfloor}^{-1}(\boldsymbol{u})\right\}\right] \tag{39}$$

and

$$\tilde{\tilde{\mathbb{Z}}}_n^{(m)}(s,\boldsymbol{u}) = \frac{1}{\sqrt{n}}\sum_{i=1}^{\lfloor ns\rfloor}\xi_i^{(m)}\left[\mathbf{1}\left\{\boldsymbol{U}_i\leq G_n^{-1}(\boldsymbol{u})\right\} - G_n\left\{G_n^{-1}(\boldsymbol{u})\right\}\right]. \tag{40}$$

The proof of Theorem 1 is based on six lemmas.

Lemma 1. *Assume that ($\mathcal{M}1$) holds. Then, for any $m\in\{1,\dots,M\}$,*

$$\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\hat{\mathbb{Z}}_n^{(m)}(s,\boldsymbol{u}) - \tilde{\tilde{\mathbb{Z}}}_n^{(m)}(s,\boldsymbol{u})\right| \xrightarrow{\mathrm{P}} 0,$$

where $\hat{\mathbb{Z}}_n^{(m)}$ and $\tilde{\tilde{\mathbb{Z}}}_n^{(m)}$ are defined in (15) and (39), respectively.

Proof. We have $\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\hat{\mathbb{Z}}_n^{(m)}(s,\boldsymbol{u})-\tilde{\tilde{\mathbb{Z}}}_n^{(m)}(s,\boldsymbol{u})\right|\leq J_n+K_n$, where

$$J_n=\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\frac{1}{\sqrt{n}}\sum_{i=1}^{\lfloor ns\rfloor}\xi_i^{(m)}\left[\mathbf{1}(\hat{\boldsymbol{U}}_{i,\lfloor ns\rfloor}\leq\boldsymbol{u})-\mathbf{1}\left\{\boldsymbol{U}_i\leq G_{\lfloor ns\rfloor}^{-1}(\boldsymbol{u})\right\}\right]\right|$$

and

$$K_n=\sup_{s\in[0,1]}\left[\left|\frac{1}{\sqrt{n}}\sum_{i=1}^{\lfloor ns\rfloor}\xi_i^{(m)}\right|\sup_{\boldsymbol{u}\in[0,1]^d}\left|C_{\lfloor ns\rfloor}(\boldsymbol{u})-G_{\lfloor ns\rfloor}\left\{G_{\lfloor ns\rfloor}^{-1}(\boldsymbol{u})\right\}\right|\right].$$

Now,

$$K_n=\max_{1\leq k\leq n}\left[\left|\frac{1}{\sqrt{n}}\sum_{i=1}^{k}\xi_i^{(m)}\right|\sup_{\boldsymbol{u}\in[0,1]^d}\left|C_k(\boldsymbol{u})-G_k\left\{G_k^{-1}(\boldsymbol{u})\right\}\right|\right]$$

and

$$\left|C_k(\boldsymbol{u})-G_k\left\{G_k^{-1}(\boldsymbol{u})\right\}\right|\leq\frac{1}{k}\sum_{i=1}^{k}\left|\mathbf{1}(\hat{\boldsymbol{U}}_{i,k}\leq\boldsymbol{u})-\mathbf{1}\left\{\boldsymbol{U}_i\leq G_k^{-1}(\boldsymbol{u})\right\}\right|,\qquad\boldsymbol{u}\in[0,1]^d.$$

From (37), we have that, for any $j\in\{1,\dots,d\}$ and $u\in[0,1]$, $U_{ij}\leq G_{k,j}^{-1}(u)$ is equivalent to $(R_{ij,k}-1)/k<u$. Since $R_{ij,k}/(k+1)-(R_{ij,k}-1)/k>0$, it follows that, for any $\boldsymbol{u}\in[0,1]^d$,

$$\begin{aligned}\left|\mathbf{1}(\hat{\boldsymbol{U}}_{i,k}\leq\boldsymbol{u})-\mathbf{1}\left\{\boldsymbol{U}_i\leq G_k^{-1}(\boldsymbol{u})\right\}\right|&=\left|\prod_{j=1}^{d}\mathbf{1}\left(\frac{R_{ij,k}}{k+1}\leq u_j\right)-\prod_{j=1}^{d}\mathbf{1}\left(\frac{R_{ij,k}-1}{k}<u_j\right)\right|\\&\leq\sum_{j=1}^{d}\mathbf{1}\left(\frac{R_{ij,k}-1}{k}<u_j<\frac{R_{ij,k}}{k+1}\right).\end{aligned}\tag{41}$$

Consequently,

$$\sup_{\boldsymbol{u}\in[0,1]^d}\left|C_k(\boldsymbol{u})-G_k\left\{G_k^{-1}(\boldsymbol{u})\right\}\right|\leq\sum_{j=1}^{d}\sup_{u_j\in[0,1]}\frac{1}{k}\sum_{i=1}^{k}\mathbf{1}\left(\frac{R_{ij,k}-1}{k}<u_j<\frac{R_{ij,k}}{k+1}\right)\leq\frac{d}{k}.\tag{42}$$

Hence,

$$K_n\leq\frac{d}{\sqrt{n}}\max_{1\leq k\leq n}\left|\frac{1}{k}\sum_{i=1}^{k}\xi_i^{(m)}\right|\leq\frac{d}{\sqrt{n}}\sup_{k\geq1}\left|\frac{1}{k}\sum_{i=1}^{k}\xi_i^{(m)}\right|\xrightarrow{\text{a.s.}}0,$$

since, from the strong law of large numbers, $\sup_{k\geq1}\left|k^{-1}\sum_{i=1}^{k}\xi_i^{(m)}\right|<\infty$ almost surely.

It remains to show that $J_n \overset{\mathrm{P}}{\to} 0$. Using (41), we obtain that

$$\begin{aligned}
J_n &\le \max_{1\le k\le n} \sup_{\boldsymbol{u}\in[0,1]^d} \frac{1}{\sqrt{n}} \sum_{i=1}^{k} \left|\xi_i^{(m)}\right| \left|\mathbf{1}(\hat{\boldsymbol{U}}_{i,k} \le \boldsymbol{u}) - \mathbf{1}\left\{\boldsymbol{U}_i \le G_k^{-1}(\boldsymbol{u})\right\}\right| \\
&\le \max_{1\le k\le n} \sup_{\boldsymbol{u}\in[0,1]^d} \frac{1}{\sqrt{n}} \sum_{i=1}^{k} \left|\xi_i^{(m)}\right| \sum_{j=1}^{d} \mathbf{1}\left(\frac{R_{ij,k}-1}{k} < u_j < \frac{R_{ij,k}}{k+1}\right) \\
&\le \frac{1}{\sqrt{n}} \sum_{j=1}^{d} \max_{1\le k\le n} \sup_{u_j\in[0,1]} \sum_{i=1}^{k} \left|\xi_i^{(m)}\right| \mathbf{1}\left(\frac{R_{ij,k}-1}{k} < u_j < \frac{R_{ij,k}}{k+1}\right) \\
&\le \frac{1}{\sqrt{n}} \sum_{j=1}^{d} \max_{1\le k\le n} \max_{1\le i\le k} \left|\xi_i^{(m)}\right| \le \frac{d}{\sqrt{n}} \max_{1\le k\le n} \left|\xi_k^{(m)}\right|.
\end{aligned}$$

The expectation of $n^{-1/2}\max_{1\le k\le n}|\xi_k^{(m)}|$ is known to converge to zero provided $\mathrm{E}\{(\xi_k^{(m)})^2\} < \infty$, which is the case since $\int_0^\infty \{\mathrm{P}(|\xi_i^{(m)}| > x)\}^{1/2}\mathrm{d}x < \infty$ (see e.g. Kosorok, 2008, Exercises 10.5.1 and 10.5.2). It follows that $J_n \overset{\mathrm{P}}{\to} 0$, which completes the proof. ■

Lemma 2. *Assume that ($\mathcal{M}2$) holds. Then, for any $m \in \{1,\dots,M\}$,*

$$\sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} \left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \hat{\mathbb{Z}}_n^{(m)}(s,\boldsymbol{u})\right| \overset{\mathrm{P}}{\to} 0,$$

where $\mathbb{Z}_n^{(m)}$ and $\hat{\mathbb{Z}}_n^{(m)}$ are defined in (14) and (15), respectively.

Proof. We have $\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \hat{\mathbb{Z}}_n^{(m)}(s,\boldsymbol{u})\right| \le J_n + K_n$, where

$$J_n = \sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} \left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \tilde{\hat{\mathbb{Z}}}_n^{(m)}(s,\boldsymbol{u})\right|$$

and

$$K_n = \sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} \left|\tilde{\hat{\mathbb{Z}}}_n^{(m)}(s,\boldsymbol{u}) - \hat{\mathbb{Z}}_n^{(m)}(s,\boldsymbol{u})\right|.$$

We know that $K_n \overset{\mathrm{P}}{\to} 0$ from Lemma 1. It remains to show that $J_n \overset{\mathrm{P}}{\to} 0$.

It can be verified that $J_n \le J_n' + J_n''$, where

$$J_n' = \sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} \left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \mathbb{Z}_n^{(m)}\left\{s, G_{\lfloor ns\rfloor}^{-1}(\boldsymbol{u})\right\}\right|,$$

and

$$J_n'' = \sup_{s\in[0,1]} \left[\left|\frac{1}{\sqrt{n}}\sum_{i=1}^{\lfloor ns\rfloor} \xi_i^{(m)}\right| \sup_{\boldsymbol{u}\in[0,1]^d} \left|G_{\lfloor ns\rfloor}\left\{G_{\lfloor ns\rfloor}^{-1}(\boldsymbol{u})\right\} - C\left\{G_{\lfloor ns\rfloor}^{-1}(\boldsymbol{u})\right\}\right|\right].$$

Now,

$$J_n'' \le \max_{1\le k\le n} \left[\left|\frac{1}{\sqrt{n}}\sum_{i=1}^{k} \xi_i^{(m)}\right| \sup_{\boldsymbol{u}\in[0,1]^d} |G_k(\boldsymbol{u}) - C(\boldsymbol{u})|\right].$$

Using the law of the iterated logarithm for i.i.d. mean 0 variance 1 sequences and the law of the iterated logarithm for multivariate c.d.f.s (see e.g. Kiefer, 1961; Dudley and Philipp, 1983), J_n'' converges to zero almost surely since

$$\max_{3\le k\le n}\left[\left|\frac{1}{\sqrt{n}}\sum_{i=1}^{k}\xi_i^{(m)}\right|\sup_{\boldsymbol{u}\in[0,1]^d}|G_k(\boldsymbol{u})-C(\boldsymbol{u})|\right]$$

$$\le \max_{3\le k\le n}\frac{\left|\sum_{i=1}^{k}\xi_i^{(m)}\right|}{(k\log\log k)^{1/2}}\times\max_{3\le k\le n}\frac{\sup_{\boldsymbol{u}\in[0,1]^d}|G_k(\boldsymbol{u})-C(\boldsymbol{u})|}{k^{-1/2}(\log\log k)^{1/2}}\times\frac{1}{\sqrt{n}}\max_{3\le k\le n}\log\log k$$

converges to zero almost surely.

Concerning J_n', we have

$$J_n' = \max_{1\le k\le n}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_n^{(m)}(k/n,\boldsymbol{u})-\mathbb{Z}_n^{(m)}\left\{k/n,G_k^{-1}(\boldsymbol{u})\right\}\right|.$$

Let $a_n = n^{-1/2}(\log\log n)^{1/2}$, $n\ge 3$. From the law of the iterated logarithm for univariate c.d.f.s (see e.g. Chung, 1949, Theorem 2*), we have that, for any $j\in\{1,\dots,d\}$,

$$\limsup\frac{\sup_{u\in[0,1]}|G_{n,j}(u)-u|}{a_n}=\frac{1}{\sqrt{2}},\qquad\text{almost surely.}\tag{43}$$

By considering the graphs of $G_{n,j}$ and $G_{n,j}^{-1}$, it immediately appears, as in Shorack and Welner (1986, page 86), that

$$\sup_{u\in[0,1]}|G_{n,j}(u)-u| = \sup_{u\in[0,1]}|G_{n,j}^{-1}(u)-u|,\tag{44}$$

from which we obtain that, for almost every ω, there exists k_ω such $k\ge k_\omega$ implies

$$\max_{1\le j\le d}\sup_{u\in[0,1]}\left|G_{k,j}^{-1}(u)(\omega)-u\right|\le a_k.$$

Hence, for almost every ω,

$$J_n'(\omega)\le\max_{1\le k<k_\omega}Y_{n,k}(\omega)+\max_{k_\omega\le k\le n}Y_{n,k}(\omega),$$

where $Y_{n,k}=\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_n^{(m)}(k/n,\boldsymbol{u})-\mathbb{Z}_n^{(m)}\left\{k/n,G_k^{-1}(\boldsymbol{u})\right\}\right|$. Now,

$$\max_{1\le k<k_\omega}Y_{n,k}(\omega)\le\frac{1}{\sqrt{n}}\sum_{i=1}^{k_\omega-1}4|\xi_i^{(m)}(\omega)|\to 0.$$

For $k\ge k_\omega$, we have

$$Y_{n,k}(\omega)\le\sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in[0,1]^d\\ \forall j\,|u_j-v_j|\le a_k}}\left|\mathbb{Z}_n^{(m)}(k/n,\boldsymbol{u})(\omega)-\mathbb{Z}_n^{(m)}(k/n,\boldsymbol{v})(\omega)\right|.$$

In other words, with probability one, J_n' is smaller than

$$L_n=\max_{1\le k\le n}\sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in[0,1]^d\\ \forall j\,|u_j-v_j|\le a_k}}\left|\mathbb{Z}_n^{(m)}(k/n,\boldsymbol{u})-\mathbb{Z}_n^{(m)}(k/n,\boldsymbol{v})\right|$$

plus a term that converges to zero almost surely.

To conclude the proof, it remains to show that $L_n \xrightarrow{\mathrm{P}} 0$. Let

$$\tilde{M}_n(a) = \sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in[0,1]^d \\ \forall j\ |u_j-v_j|\le a}} \left| \frac{1}{\sqrt{n}} \sum_{i=1}^n \xi_i^{(m)} Z_i(\boldsymbol{u},\boldsymbol{v}) \right| = \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n \xi_i^{(m)} Z_i \right\|_a, \qquad a \ge 0,$$

where $Z_i(\boldsymbol{u},\boldsymbol{v}) = \mathbf{1}(\boldsymbol{U}_i \le \boldsymbol{u}) - C(\boldsymbol{u}) - \mathbf{1}(\boldsymbol{U}_i \le \boldsymbol{v}) + C(\boldsymbol{v})$ and, for any $f : [0,1]^{2d} \to \mathbb{R}$, $\|f\|_a = \sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in[0,1]^d \\ \forall j\ |u_j-v_j|\le a}} |f(\boldsymbol{u},\boldsymbol{v})|$. Then, L_n can be rewritten as

$$L_n = \max_{1\le k\le n} \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^k \xi_i^{(m)} Z_i \right\|_{a_k} = \max_{1\le k\le n} \sqrt{\frac{k}{n}} \tilde{M}_k(a_k).$$

Let $\nu \ge 1$. Then,

$$\begin{aligned} \mathrm{E}\left(L_n^\nu\right) = \mathrm{E}\left[\max_{1\le k\le n} \left(\frac{k}{n}\right)^{\nu/2} \left\{\tilde{M}_k(a_k)\right\}^\nu \right] &\le \mathrm{E}\left[\sum_{k=1}^n \left(\frac{k}{n}\right)^{\nu/2} \left\{\tilde{M}_k(a_k)\right\}^\nu \right] \\ &\le n \max_{1\le k\le n} \left(\frac{k}{n}\right)^{\nu/2} \mathrm{E}\left[\left\{\tilde{M}_k(a_k)\right\}^\nu\right]. \end{aligned}$$

From Lemma 2.3.6 of van der Vaart and Wellner (2000) with $\Phi(x) = x^\nu$, $x \ge 0$, we obtain that, for any $k \ge 1$,

$$\mathrm{E}\left[\left\{\tilde{M}_k(a_k)\right\}^\nu\right] = \mathrm{E}\left[\left\| \frac{1}{\sqrt{k}} \sum_{i=1}^k \xi_i^{(m)} Z_i \right\|_{a_k}^\nu \right] \le 2^\nu \mathrm{E}\left[\left\| \frac{1}{\sqrt{k}} \sum_{i=1}^k Z_i \right\|_{a_k}^\nu \right].$$

It follows that

$$\mathrm{E}\left(L_n^\nu\right) \le 2^\nu n \max_{1\le k\le n} \left(\frac{k}{n}\right)^{\nu/2} \mathrm{E}\left[\{M_k(a_k)\}^\nu\right],$$

where

$$M_n(a) = \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n Z_i \right\|_a, \qquad a \ge 0,$$

is the oscillation modulus studied by Einmahl (1987). From his Theorem 5.2, we have that, for sufficiently large n,

$$\mathrm{E}\left[\{M_n(a)\}^\nu\right] = O\left((a \log a^{-1})^{\nu/2}\right)$$

when C is the independence copula. The proof of that result is based on Inequality 5.3 on page 73 of Einmahl (1987). As explained in Segers (2012, Appendix A), this inequality continues to hold for any copula C. Consequently, we have that

$$\mathrm{E}\left[\{M_n(a_n)\}^\nu\right] = O\left(b_n\right),$$

where $b_n = n^{-\nu/4} (\log\log n)^{\nu/4} (\log n)^{\nu/2}$.

Then, we write

$$2^\nu n \max_{3\leq k\leq n}\left(\frac{k}{n}\right)^{\nu/2} \mathrm{E}\left[\{M_k(a_k)\}^\nu\right] \leq 2^\nu n \max_{3\leq k\leq n}\left(\frac{k}{n}\right)^{\nu/2} b_k \times \max_{3\leq k\leq n}\frac{\mathrm{E}\left[\{M_k(a_k)\}^\nu\right]}{b_k}.$$

The last maximum is bounded, while

$$2^\nu n \max_{3\leq k\leq n}\left(\frac{k}{n}\right)^{\nu/2} b_k = 2^\nu n^{1-\nu/2}\max_{3\leq k\leq n} k^{\nu/2} b_k = 2^\nu n^{1-\nu/4}(\log\log n)^{\nu/4}(\log n)^{\nu/2},$$

which converges to zero for $\nu > 4$. It follows that $\mathrm{E}\left(L_n^\nu\right)$ converges to zero for any $\nu > 4$, which implies that $L_n \overset{\mathrm{P}}{\to} 0$. ■

Lemma 3. *Assume that ($\mathcal{M}2$) holds. Then, for any $m \in \{1,\dots,M\}$ and $j \in \{1,\dots,d\}$,*

$$\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d}\left[\left|C^{[j]}_{\lfloor ns\rfloor}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{u})\right|\left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}^{(j)})\right|\right] \overset{\mathrm{P}}{\to} 0,$$

where $C^{[j]}_{\lfloor ns\rfloor}$ is defined analogously to (17), and $\mathbb{Z}_n^{(m)}$ is defined in (14).

Proof. For any $\delta \in (0,1/2)$, we have

$$\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d}\left[\left|C^{[j]}_{\lfloor ns\rfloor}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{u})\right|\left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}^{(j)})\right|\right] \leq J_n + K_n,$$

where

$$J_n = \max_{1\leq k\leq n}\left[\frac{k^{1/4}}{n^{1/8}}\sup_{\substack{\boldsymbol{u}\in[0,1]^d\\ u_j\in[\delta,1-\delta]}}\left|C^{[j]}_k(\boldsymbol{u}) - C^{[j]}(\boldsymbol{u})\right|\frac{k^{1/4}}{n^{3/8}}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_k^{(m)}(1,\boldsymbol{u}^{(j)})\right|\right]$$

and

$$K_n = 5\sup_{s\in[0,1]}\sup_{\substack{\boldsymbol{u}\in[0,1]^d\\ u_j\notin[\delta,1-\delta]}}\left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}^{(j)})\right|,$$

since $C^{[j]}_n(\boldsymbol{u}) \leq 5$ for all $\boldsymbol{u} \in [0,1]^d$ and $n \geq 1$ as shown in Kojadinovic et al. (2011, Proof of Proposition 2). Now, since, for any $s \in [0,1]$, $\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) = 0$ when $\boldsymbol{u}$ is on the boundary of $[0,1]^d$,

$$\begin{aligned}\sup_{s\in[0,1]}\sup_{\substack{\boldsymbol{u}\in[0,1]^d\\ u_j\notin[\delta,1-\delta]}}\left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}^{(j)})\right| &\leq \sup_{s\in[0,1]}\sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in[0,1]^d\\ |u_j-v_j|<\delta}}\left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}^{(j)}) - \mathbb{Z}_n^{(m)}(s,\boldsymbol{v}^{(j)})\right|\\ &\leq \sup_{\substack{s,t\in[0,1],\boldsymbol{u},\boldsymbol{v}\in[0,1]^d\\ |s-t|+\rho(u_j,v_j)\leq\sqrt{\delta+\delta^2}}}\left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}^{(j)}) - \mathbb{Z}_n^{(m)}(t,\boldsymbol{v}^{(j)})\right|,\end{aligned}$$

where

$$\rho^2(u,v) = \mathrm{E}\left[\{1(U\leq u) - u - 1(U\leq v) + v\}^2\right] = |u-v| - (u-v)^2,$$

and U is a standard uniform random variable. The weak convergence of $\mathbb{Z}_n^{(m)}$ implies the asymptotic equicontinuity of $(s, u_j) \mapsto \mathbb{Z}_n^{(m)}(s, 1, \dots, 1, u_j, 1, \dots, 1)$ with respect to the natural semimetric on $[0,1] \times [0,1]$ (see van der Vaart and Wellner, 2000, Chapter 2.12), i.e., for any $\varepsilon > 0$,

$$\lim_{\gamma \downarrow 0} \limsup_{n \to \infty} \mathrm{P}\left\{ \sup_{\substack{s,t \in [0,1], \boldsymbol{u}, \boldsymbol{v} \in [0,1]^d \\ |s-t| + \rho(u_j, v_j) \leq \gamma}} \left| \mathbb{Z}_n^{(m)}(s, \boldsymbol{u}^{(j)}) - \mathbb{Z}_n^{(m)}(t, \boldsymbol{v}^{(j)}) \right| > \varepsilon \right\} = 0.$$

Hence, for fixed $\varepsilon > 0$ and $\eta > 0$, we can choose $\delta > 0$ sufficiently small so that, for sufficiently large n, $\mathrm{P}(K_n > \varepsilon) < \eta/2$.

Let us now show that, for the chosen δ, J_n converges in probability to zero, which will imply that, for sufficiently large n, $\mathrm{P}(J_n > \varepsilon) < \eta/2$ and will complete the proof.

From the law of the iterated logarithm stated in (43), we have that, for any $j \in \{1, \dots, d\}$,

$$\sup_{n \geq 3} \frac{\sup_{u \in [0,1]} n^{1/2} \left| G_{n,j}(u) - u \right|}{(\log \log n)^{1/2}} < \infty \qquad \text{almost surely.}$$

Since $|\mathbf{1}(U_{nj} \leq u) - u| \leq 1$ for all $u \in [0,1]$ and $n \geq 1$, for any $\nu > 0$,

$$\mathrm{E}\left[\sup_{n \geq 3} \frac{\left\{ \sup_{u \in [0,1]} \left| \mathbf{1}(U_{nj} \leq u) - u \right| \right\}^\nu}{(n \log \log n)^{\nu/2}} \right] < \infty,$$

which, from Corollary A.1.8 of van der Vaart and Wellner (2000), is equivalent to

$$\mathrm{E}\left[\sup_{n \geq 3} \frac{\left\{ \sup_{u \in [0,1]} n^{1/2} \left| G_{n,j}(u) - u \right| \right\}^\nu}{(\log \log n)^{\nu/2}} \right] < \infty,$$

and implies that

$$\sup_{n \geq 3} \frac{\mathrm{E}\left[\left\{ \sup_{u \in [0,1]} n^{1/2} \left| G_{n,j}(u) - u \right| \right\}^\nu \right]}{(\log \log n)^{\nu/2}} \leq \mathrm{E}\left[\sup_{n \geq 3} \frac{\left\{ \sup_{u \in [0,1]} n^{1/2} \left| G_{n,j}(u) - u \right| \right\}^\nu}{(\log \log n)^{\nu/2}} \right] < \infty.$$

Let $\nu \geq 1$. Applying Lemma 2.3.6 of van der Vaart and Wellner (2000) with $\Phi(x) = x^\nu$, $x \geq 0$, we obtain

$$\sup_{n \geq 3} \mathrm{E}\left[\frac{\left\{ \sup_{\boldsymbol{u} \in [0,1]^d} \left| \mathbb{Z}_n^{(m)}(1, \boldsymbol{u}^{(j)}) \right| \right\}^\nu}{(\log \log n)^{\nu/2}} \right] \leq \sup_{n \geq 3} \mathrm{E}\left[2^\nu \frac{\left\{ \sup_{u \in [0,1]} n^{1/2} \left| G_{n,j}(u) - u \right| \right\}^\nu}{(\log \log n)^{\nu/2}} \right] < \infty. \tag{45}$$

Let us now show that $J_n \xrightarrow{\mathrm{P}} 0$. We have

$$\max_{3 \leq k \leq n} \left[\frac{k^{1/4}}{n^{1/8}} \sup_{\substack{\boldsymbol{u} \in [0,1]^d \\ u_j \in [\delta, 1-\delta]}} \left| C_k^{[j]}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{u}) \right| \frac{k^{1/4}}{n^{3/8}} \sup_{\boldsymbol{u} \in [0,1]^d} \left| \mathbb{Z}_k^{(m)}(1, \boldsymbol{u}^{(j)}) \right| \right] \leq L_n \times M_n,$$

where

$$L_n = \max_{3\le k\le n} \sup_{\substack{\boldsymbol{u}\in[0,1]^d \\ u_j\in[\delta,1-\delta]}} \frac{\left|C_k^{[j]}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{u})\right|}{k^{-1/4}(\log k)^{1/2}(\log\log k)^{1/4}} \times n^{-1/8} \max_{3\le k\le n}(\log k)^{1/2}(\log\log k)^{3/4}$$

and

$$M_n = \max_{3\le k\le n}\left[\frac{k^{1/4}}{n^{3/8}}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_k^{(m)}(1,\boldsymbol{u}^{(j)})\right|\right].$$

The first maximum in L_n is strictly smaller than ∞ almost surely by Proposition 2, which implies that $L_n \xrightarrow{\text{a.s.}} 0$. It remains to show that $M_n \overset{\text{P}}{\to} 0$. For $\nu \ge 1$, we have

$$\begin{aligned}
\mathrm{E}\left(M_n^\nu\right) &= \mathrm{E}\max_{3\le k\le n}\left[\frac{k^{\nu/4}}{n^{3\nu/8}}\left\{\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_k^{(m)}(1,\boldsymbol{u}^{(j)})\right|\right\}^\nu\right] \\
&\le \mathrm{E}\left[\sum_{k=3}^n \frac{k^{\nu/4}}{n^{3\nu/8}}\left\{\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_k^{(m)}(1,\boldsymbol{u}^{(j)})\right|\right\}^\nu\right] \\
&\le n\max_{3\le k\le n}\frac{k^{\nu/4}}{n^{3\nu/8}}\mathrm{E}\left[\left\{\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_k^{(m)}(1,\boldsymbol{u}^{(j)})\right|\right\}^\nu\right] \\
&\le n^{1-3\nu/8}\max_{3\le k\le n}k^{\nu/4}(\log\log k)^{\nu/2} \times \max_{3\le k\le n}\frac{\mathrm{E}\left[\left\{\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_k^{(m)}(1,\boldsymbol{u}^{(j)})\right|\right\}^\nu\right]}{(\log\log k)^{\nu/2}}.
\end{aligned}$$

The second maximum on the right of the last inequality is bounded by (45), while the first maximum is equal to $n^{1-\nu/8}(\log\log n)^{\nu/2}$, which converges to zero for $\nu > 8$. Hence, $M_n \overset{\text{P}}{\to} 0$, which completes the proof. ■

Lemma 4. *Assume that ($\mathcal{M}1$) holds. Then, for any $m \in \{1,\dots,M\}$,*

$$\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\check{\mathbb{Z}}_n^{(m)}(s,\boldsymbol{u}) - \tilde{\check{\mathbb{Z}}}_n^{(m)}(s,\boldsymbol{u})\right| \overset{\text{P}}{\to} 0,$$

where $\check{\mathbb{Z}}_n^{(m)}$ and $\tilde{\check{\mathbb{Z}}}_n^{(m)}$ are defined in (16) and (40), respectively.

Proof. The proof is a simpler version of the proof of Lemma 1 and is omitted. ■

Lemma 5. *Assume that ($\mathcal{M}1$) holds. Then, for any $m \in \{1,\dots,M\}$,*

$$\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \check{\mathbb{Z}}_n^{(m)}(s,\boldsymbol{u})\right| \overset{\text{P}}{\to} 0,$$

where $\mathbb{Z}_n^{(m)}$ and $\check{\mathbb{Z}}_n^{(m)}$ are defined in (14) and (16), respectively.

Proof. We have $\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \check{\mathbb{Z}}_n^{(m)}(s,\boldsymbol{u})\right| \le J_n + K_n$, where

$$J_n = \sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \tilde{\check{\mathbb{Z}}}_n^{(m)}(s,\boldsymbol{u})\right|$$

and

$$K_n = \sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} \left| \tilde{\check{\mathbb{Z}}}_n^{(m)}(s,\boldsymbol{u}) - \check{\mathbb{Z}}_n^{(m)}(s,\boldsymbol{u}) \right|.$$

We know that $K_n \overset{\mathrm{P}}{\to} 0$ from Lemma 4. It remains to show that $J_n \overset{\mathrm{P}}{\to} 0$.

It can be verified that $J_n \leq J_n' + J_n''$, where

$$J_n' = \sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} \left| \mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \mathbb{Z}_n^{(m)} \left\{ s, G_n^{-1}(\boldsymbol{u}) \right\} \right|$$

and

$$J_n'' = \sup_{s\in[0,1]} \left| \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} \xi_i^{(m)} \right| \times \sup_{\boldsymbol{u}\in[0,1]^d} \left| G_n\left\{ G_n^{-1}(\boldsymbol{u}) \right\} - C\left\{ G_n^{-1}(\boldsymbol{u}) \right\} \right|.$$

Now,

$$J_n'' \leq \sup_{s\in[0,1]} \left| \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} \xi_i^{(m)} \right| \times \sup_{\boldsymbol{u}\in[0,1]^d} \left| G_n(\boldsymbol{u}) - C(\boldsymbol{u}) \right| \overset{\mathrm{P}}{\to} 0$$

since the first supremum converges weakly to the supremum of a Brownian motion and the second supremum converges almost surely to zero by the Glivenko-Cantelli lemma.

Concerning J_n', we have, using (43), that, for n sufficiently large,

$$J_n' \leq \sup_{s\in[0,1]} \sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in[0,1]^d \\ \forall j\ |u_j - v_j| \leq a_n}} \left| \mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \mathbb{Z}_n^{(m)}(s,\boldsymbol{v}) \right|,$$

where $a_n = n^{-1/2}(\log\log n)^{1/2}$, $n \geq 3$. The quantity on the right converges in probability to zero due to the asymptotic equicontinuity of $\mathbb{Z}_n^{(m)}$. Indeed, as discussed in van der Vaart and Wellner (2000, Chapter 2.12), the asymptotic equicontinuity of $\mathbb{Z}_n^{(m)}$ can be stated in terms of the natural semimetric on $[0,1] \times [0,1]^d$, for any decreasing sequence $\delta_n \downarrow 0$, as

$$\sup_{\substack{s,t\in[0,1], \boldsymbol{u},\boldsymbol{v}\in[0,1]^d \\ |s-t| + \rho_C(\boldsymbol{u},\boldsymbol{v}) \leq \delta_n}} \left| \mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \mathbb{Z}_n^{(m)}(t,\boldsymbol{v}) \right| \overset{\mathrm{P}}{\to} 0,$$

where $\rho_C^2(\boldsymbol{u},\boldsymbol{v}) = \mathrm{E}\left[\{ 1(\boldsymbol{U} \leq \boldsymbol{u}) - C(\boldsymbol{u}) - 1(\boldsymbol{U} \leq \boldsymbol{v}) + C(\boldsymbol{v}) \}^2 \right]$, and $\boldsymbol{U}$ is a random vector with c.d.f. C.

Now, it is easy to verify that

$$\rho_C^2(\boldsymbol{u},\boldsymbol{v}) = \left| C(\boldsymbol{u}) - C(\boldsymbol{u} \wedge \boldsymbol{v}) + C(\boldsymbol{v}) - C(\boldsymbol{u} \wedge \boldsymbol{v}) - \{ C(\boldsymbol{u}) - C(\boldsymbol{v}) \}^2 \right|,$$

which implies that

$$\rho_C^2(\boldsymbol{u},\boldsymbol{v}) \leq C(\boldsymbol{u} \vee \boldsymbol{v}) - C(\boldsymbol{u} \wedge \boldsymbol{v}) + \{ C(\boldsymbol{u}) - C(\boldsymbol{v}) \}^2 \\ \leq d \max_{1\leq j\leq d} |u_j - v_j| + d^2 \max_{1\leq j\leq d} |u_j - v_j|^2,$$

since C satisfies the Lipschitz condition

$$|C(\boldsymbol{u}) - C(\boldsymbol{v})| \leq \sum_{j=1}^{d} |u_j - v_j|, \qquad \boldsymbol{u},\boldsymbol{v} \in [0,1]^d. \tag{46}$$

Hence, with $\delta_n = \sqrt{da_n + d^2 a_n^2}$,

$$J_n' \leq \sup_{s\in[0,1]} \sup_{\substack{\boldsymbol{u},\boldsymbol{v}\in[0,1]^d \\ \forall j\,|u_j - v_j|\leq a_n}} \left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \mathbb{Z}_n^{(m)}(s,\boldsymbol{v})\right| \leq \sup_{\substack{s\in[0,1],\boldsymbol{u},\boldsymbol{v}\in[0,1]^d \\ \rho_C(\boldsymbol{u},\boldsymbol{v})\leq\delta_n}} \left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \mathbb{Z}_n^{(m)}(s,\boldsymbol{v})\right|$$

$$\leq \sup_{\substack{s,t\in[0,1],\boldsymbol{u},\boldsymbol{v}\in[0,1]^d \\ |s-t|+\rho_C(\boldsymbol{u},\boldsymbol{v})\leq\delta_n}} \left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \mathbb{Z}_n^{(m)}(t,\boldsymbol{v})\right| \xrightarrow{\mathrm{P}} 0,$$

which completes the proof. ■

Lemma 6. *Assume that ($\mathcal{M}1$) holds. Then, for any $m \in \{1,\dots,M\}$ and $j \in \{1,\dots,d\}$,*

$$\sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} \left[\left|C_n^{[j]}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{u})\right| \left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}^{(j)})\right|\right] \xrightarrow{\mathrm{P}} 0,$$

where $C_n^{[j]}$ and $\mathbb{Z}_n^{(m)}$ are defined in (17) and (14), respectively.

Proof. For any $\delta \in (0,1/2)$, we have

$$\sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} \left[\left|C_n^{[j]}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{u})\right| \left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}^{(j)})\right|\right] \leq J_n + K_n,$$

where

$$J_n = \sup_{\substack{\boldsymbol{u}\in[0,1]^d \\ u_j\in[\delta,1-\delta]}} \left|C_n^{[j]}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{u})\right| \sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} \left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}^{(j)})\right|$$

and

$$K_n = 5 \sup_{s\in[0,1]} \sup_{\substack{\boldsymbol{u}\in[0,1]^d \\ u_j\notin[\delta,1-\delta]}} \left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}^{(j)})\right|,$$

since $C_n^{[j]}(\boldsymbol{u}) \leq 5$ for all $\boldsymbol{u} \in [0,1]^d$ and $n \geq 1$. Proceeding as in the proof of Lemma 3, for fixed $\varepsilon > 0$ and $\eta > 0$, we can choose $\delta > 0$ sufficiently small so that, for sufficiently large n, $\mathrm{P}(K_n > \varepsilon) < \eta/2$. For that δ, J_n converges in probability to zero by Proposition 2 and since $\sup_{s\in[0,1],\boldsymbol{u}\in[0,1]^d} \left|\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}^{(j)})\right| = O_{\mathrm{P}}(1)$. It follows that, for sufficiently large n, $\mathrm{P}(J_n > \varepsilon) < \eta/2$, which completes the proof. ■

Proof of Theorem 1. We only prove (ii) as the proof of (i) is similar. A consequence of the sequential extension of the multiplier central limit theorem proved in Holmes et al. (2013, Theorem 1) is that

$$\left(\mathbb{Z}_n, \mathbb{Z}_n^{(1)}, \dots, \mathbb{Z}_n^{(M)}\right) \rightsquigarrow \left(\mathbb{Z}_C, \mathbb{Z}_C^{(1)}, \dots, \mathbb{Z}_C^{(M)}\right) \tag{47}$$

in $\{\ell^\infty([0,1]^{d+1})\}^{M+1}$, where $\mathbb{Z}_n$ and $\mathbb{Z}_n^{(m)}$ are defined in (9) and (14), respectively, $\mathbb{Z}_C$ is a *C-Kiefer-Müller* process and $\mathbb{Z}_C^{(1)}, \dots, \mathbb{Z}_C^{(M)}$ are independent copies of $\mathbb{Z}_C$.

Now, for any $m \in \{1,\dots,M\}$, let

$$\mathbb{C}_n^{(m)}(s,\boldsymbol{u}) = \mathbb{Z}_n^{(m)}(s,\boldsymbol{u}) - \sum_{j=1}^d C^{[j]}(\boldsymbol{u})\mathbb{Z}_n^{(m)}(s,\boldsymbol{u}^{(j)}), \qquad (s,\boldsymbol{u}) \in [0,1]^{d+1}.$$

Then, from the continuous mapping theorem and the fact that A_n defined in (34) converges in probability to zero, we have that

$$\left(\mathbb{C}_n, \mathbb{C}_n^{(1)}, \dots, \mathbb{C}_n^{(M)}, \mathbb{C}_n^{(1)}, \dots, \mathbb{C}_n^{(M)}\right) \leadsto \left(\mathbb{C}_C, \mathbb{C}_C^{(1)}, \dots, \mathbb{C}_C^{(M)}, \mathbb{C}_C^{(1)}, \dots, \mathbb{C}_C^{(M)}\right) \tag{48}$$

in $\{\ell^\infty([0,1]^{d+1})\}^{2M+1}$, where $\mathbb{C}_C$ is defined in (13), and $\mathbb{C}_C^{(1)}, \dots, \mathbb{C}_C^{(M)}$ are independent copies of $\mathbb{C}_C$. Next, from Lemmas 3 and 6, we obtain that

$$\left(\mathbb{C}_n, \tilde{\hat{\mathbb{C}}}_n^{(1)}, \dots, \tilde{\hat{\mathbb{C}}}_n^{(M)}, \tilde{\check{\mathbb{C}}}_n^{(1)}, \dots, \tilde{\check{\mathbb{C}}}_n^{(M)}\right) \leadsto \left(\mathbb{C}_C, \mathbb{C}_C^{(1)}, \dots, \mathbb{C}_C^{(M)}, \mathbb{C}_C^{(1)}, \dots, \mathbb{C}_C^{(M)}\right)$$

in $\{\ell^\infty([0,1]^{d+1})\}^{2M+1}$, where, for any $m \in \{1, \dots, M\}$ and $(s, \boldsymbol{u}) \in [0,1]^{d+1}$,

$$\tilde{\hat{\mathbb{C}}}_n^{(m)}(s, \boldsymbol{u}) = \mathbb{Z}_n^{(m)}(s, \boldsymbol{u}) - \sum_{j=1}^{d} C_{\lfloor ns \rfloor}^{[j]}(\boldsymbol{u}) \mathbb{Z}_n^{(m)}(s, \boldsymbol{u}^{(j)}),$$

and

$$\tilde{\check{\mathbb{C}}}_n^{(m)}(s, \boldsymbol{u}) = \mathbb{Z}_n^{(m)}(s, \boldsymbol{u}) - \sum_{j=1}^{d} C_n^{[j]}(\boldsymbol{u}) \mathbb{Z}_n^{(m)}(s, \boldsymbol{u}^{(j)}).$$

The desired result finally follows from Lemmas 2 and 5, and the fact that $C_n^{[j]}(\boldsymbol{u}) \leq 5$ for all $\boldsymbol{u} \in [0,1]^d$ and $n \geq 1$. ■

C Proofs of Propositions 3, 4 and 5

Proof of Proposition 3. Under H_0, $\mathbb{D}_n$ can be rewritten as

$$\mathbb{D}_n(s, \boldsymbol{u}) = \{1 - \lambda_n(s)\} \mathbb{C}_n(s, \boldsymbol{u}) - \lambda_n(s) \mathbb{C}_n^\star(s, \boldsymbol{u}),$$

where $\mathbb{C}_n$ is defined in (3) and where, for any $(s, \boldsymbol{u}) \in [0,1]^{d+1}$,

$$\mathbb{C}_n^\star(s, \boldsymbol{u}) = \sqrt{n} \{1 - \lambda_n(s)\} \left\{C_{n-\lfloor ns \rfloor}^\star(\boldsymbol{u}) - C(\boldsymbol{u})\right\} = \frac{1}{\sqrt{n}} \sum_{i=\lfloor ns \rfloor + 1}^{n} \left\{\mathbf{1}(\hat{\boldsymbol{U}}_{i,n-\lfloor ns \rfloor}^\star \leq \boldsymbol{u}) - C(\boldsymbol{u})\right\}. \tag{49}$$

Let us first show that

$$A_n^\star = \sup_{s \in [0,1]} \sup_{\boldsymbol{u} \in [0,1]^d} \left| \mathbb{C}_n^\star(s, \boldsymbol{u}) - \frac{1}{\sqrt{n}} \sum_{i=\lfloor ns \rfloor + 1}^{n} \{\mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}) - C(\boldsymbol{u})\} \right.$$
$$\left. + \sum_{j=1}^{d} C^{[j]}(\boldsymbol{u}) \times \frac{1}{\sqrt{n}} \sum_{i=\lfloor ns \rfloor + 1}^{n} \left\{\mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}^{(j)}) - u_j\right\} \right| \xrightarrow{\mathrm{P}} 0. \tag{50}$$

Proceeding as in the proof of Proposition 1, we have that

$$A_n^\star = \max_{0 \leq k \leq n-1} \sqrt{n - k/n} B_{n-k}^\star = \max_{1 \leq k \leq n} \sqrt{k/n} B_k^\star,$$

where

$$B^\star_{n-k} = \sup_{\boldsymbol{u}\in[0,1]^d} \frac{1}{\sqrt{n-k}} \left| \sum_{i=k+1}^n \left\{ \mathbf{1}(\hat{\boldsymbol{U}}^\star_{i,n-k} \leq \boldsymbol{u}) - C(\boldsymbol{u}) \right\} - \sum_{i=k+1}^n \{ \mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}) - C(\boldsymbol{u}) \} \right.$$
$$\left. + \sum_{j=1}^d C^{[j]}(\boldsymbol{u}) \times \sum_{i=k+1}^n \left\{ \mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}^{(j)}) - u_j \right\} \right|, \qquad k \in \{0,\dots,n-1\}.$$

Now, clearly, for any $k \in \{1,\dots,n\}$, $(\hat{\boldsymbol{U}}_{1,k},\dots,\hat{\boldsymbol{U}}_{k,k})$ and $(\hat{\boldsymbol{U}}^\star_{n-k+1,k},\dots,\hat{\boldsymbol{U}}^\star_{n,k})$ have the same distribution. As a consequence, for any $k \in \{1,\dots,n\}$, $B^\star_k$ and B_k defined in (35) have the same distribution. More generally, $(B_1,\dots,B_n)$ and $(B^\star_1,\dots,B^\star_n)$ have the same distribution. Indeed, $(B^\star_1,\dots,B^\star_n)$ is nothing else than the version of $(B_1,\dots,B_n)$ computed from the sequence $\boldsymbol{X}_n, \boldsymbol{X}_{n-1},\dots,\boldsymbol{X}_1$. It follows that $A^\star_n$ and A_n defined in (34) have the same distribution. Consequently, $A_n \xrightarrow{\text{a.s.}} 0$ implies that $A^\star_n \xrightarrow{\text{P}} 0$.

Next, from the weak convergence of $\mathbb{Z}_n$ to $\mathbb{Z}_C$ in $\ell^\infty([0,1]^{d+1})$ and the continuous mapping theorem, we have that $(s,\boldsymbol{u}) \mapsto \mathbb{Z}_n(s,\boldsymbol{u}) - \sum_{j=1}^d C^{[j]}(\boldsymbol{u})\mathbb{Z}_n(s,\boldsymbol{u}^{(j)})$ and

$$(s,\boldsymbol{u}) \mapsto \frac{1}{\sqrt{n}} \sum_{i=\lfloor ns \rfloor+1}^n \{ \mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}) - C(\boldsymbol{u}) \} - \sum_{j=1}^d C^{[j]}(\boldsymbol{u}) \times \frac{1}{\sqrt{n}} \sum_{i=\lfloor ns \rfloor+1}^n \left\{ \mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}^{(j)}) - u_j \right\}$$
$$= \mathbb{Z}_n(1,\boldsymbol{u}) - \mathbb{Z}_n(s,\boldsymbol{u}) - \sum_{j=1}^d C^{[j]}(\boldsymbol{u}) \left\{ \mathbb{Z}_n(1,\boldsymbol{u}^{(j)}) - \mathbb{Z}_n(s,\boldsymbol{u}^{(j)}) \right\}.$$

jointly converge weakly to $\mathbb{C}_C$ and $\mathbb{C}^\star_C$, respectively, in $\{\ell^\infty([0,1]^{d+1})\}^2$. The desired result is finally a consequence of the fact that $A_n \xrightarrow{\text{a.s.}} 0$ and $A^\star_n \xrightarrow{\text{P}} 0$, and the continuous mapping theorem. The expression of the covariance function follows from simple calculations. ■

Proof of Proposition 4. We only prove (ii) as the proof of (i) is similar. For any $m \in \{1,\dots,M\}$ and $(s,\boldsymbol{u}) \in [0,1]^{d+1}$, let

$$\mathbb{C}^{\star,(m)}_n(s,\boldsymbol{u}) = \mathbb{Z}^{(m)}_n(1,\boldsymbol{u}) - \mathbb{Z}^{(m)}_n(s,\boldsymbol{u}) - \sum_{j=1}^d C^{[j]}(\boldsymbol{u}) \left\{ \mathbb{Z}^{(m)}_n(1,\boldsymbol{u}^{(j)}) - \mathbb{Z}^{(m)}_n(s,\boldsymbol{u}^{(j)}) \right\},$$

where $\mathbb{Z}^{(m)}_n$ is defined in (14). From (47), the continuous mapping theorem and the fact that $A^\star_n$ defined in (50) converges to zero in probability, we obtain that

$$\left(\mathbb{C}^\star_n, \mathbb{C}^{\star,(1)}_n, \dots, \mathbb{C}^{\star,(M)}_n, \mathbb{C}^{\star,(1)}_n, \dots, \mathbb{C}^{\star,(M)}_n \right) \rightsquigarrow \left(\mathbb{C}^\star_C, \mathbb{C}^{\star,(1)}_C, \dots, \mathbb{C}^{\star,(M)}_C, \mathbb{C}^{\star,(1)}_C, \dots, \mathbb{C}^{\star,(M)}_C \right) \tag{51}$$

in $\{\ell^\infty([0,1]^{d+1})\}^{2M+1}$, where $\mathbb{C}^\star_n$ is defined in (49), $\mathbb{C}^\star_C(s,\boldsymbol{u}) = \mathbb{C}_C(1,\boldsymbol{u}) - \mathbb{C}_C(s,\boldsymbol{u})$ for all $(s,\boldsymbol{u}) \in [0,1]^{d+1}$ and $\mathbb{C}^{\star,(1)}_C,\dots,\mathbb{C}^{\star,(M)}_C$ are independent copies of $\mathbb{C}^\star_C$.

Next, let us show that, for any $m \in \{1,\dots,M\}$,

$$J_n = \sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} \left[\left| C^{[j],\star}_{n-\lfloor ns \rfloor}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{u}) \right| \left| \mathbb{Z}^{(m)}_n(1,\boldsymbol{u}^{(j)}) - \mathbb{Z}^{(m)}_n(s,\boldsymbol{u}^{(j)}) \right| \right] \xrightarrow{\text{P}} 0.$$

It is easy to verify that J_n, written as a maximum over $1 \leq \lfloor ns \rfloor \leq n$, is nothing else than the version of $\sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} \left[\left| C^{[j]}_{\lfloor ns \rfloor}(\boldsymbol{u}) - C^{[j]}(\boldsymbol{u}) \right| \left| \mathbb{Z}_n^{(m)}(s, \boldsymbol{u}^{(j)}) \right| \right]$ computed from the sequence $(\xi_n^{(m)}, \boldsymbol{X}_n), (\xi_{n-1}^{(m)}, \boldsymbol{X}_{n-1}), \dots, (\xi_1^{(m)}, \boldsymbol{X}_1)$, which implies that the latter quantity and J_n have the same distribution. Lemma 3 therefore implies that $J_n \overset{\mathrm{P}}{\to} 0$. From the latter fact and Lemma 6, we therefore obtain that

$$\left(\mathbb{C}_n^\star, \tilde{\hat{\mathbb{C}}}_n^{\star,(1)}, \dots, \tilde{\hat{\mathbb{C}}}_n^{\star,(M)}, \tilde{\check{\mathbb{C}}}_n^{\star,(1)}, \dots, \tilde{\check{\mathbb{C}}}_n^{\star,(M)}\right) \rightsquigarrow \left(\mathbb{C}_C^\star, \mathbb{C}_C^{\star,(1)}, \dots, \mathbb{C}_C^{\star,(M)}, \mathbb{C}_C^{\star,(1)}, \dots, \mathbb{C}_C^{\star,(M)}\right)$$

in $\{\ell^\infty([0,1]^{d+1})\}^{2M+1}$, where, for any $m \in \{1, \dots, M\}$ and $(s, \boldsymbol{u}) \in [0,1]^{d+1}$,

$$\tilde{\hat{\mathbb{C}}}_n^{\star,(m)}(s, \boldsymbol{u}) = \mathbb{Z}_n^{(m)}(1, \boldsymbol{u}) - \mathbb{Z}_n^{(m)}(s, \boldsymbol{u}) - \sum_{j=1}^d C^{[j],\star}_{n-\lfloor ns \rfloor}(\boldsymbol{u}) \left\{ \mathbb{Z}_n^{(m)}(1, \boldsymbol{u}^{(j)}) - \mathbb{Z}_n^{(m)}(s, \boldsymbol{u}^{(j)}) \right\},$$

and

$$\tilde{\check{\mathbb{C}}}_n^{\star,(m)}(s, \boldsymbol{u}) = \mathbb{Z}_n^{(m)}(1, \boldsymbol{u}) - \mathbb{Z}_n^{(m)}(s, \boldsymbol{u}) - \sum_{j=1}^d C^{[j]}_n(\boldsymbol{u}) \left\{ \mathbb{Z}_n^{(m)}(1, \boldsymbol{u}^{(j)}) - \mathbb{Z}_n^{(m)}(s, \boldsymbol{u}^{(j)}) \right\}.$$

Now, reasoning as previously, for any $m \in \{1, \dots, M\}$,

$$K_n = \sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} \left| \mathbb{Z}_n^{(m)}(1, \boldsymbol{u}) - \mathbb{Z}_n^{(m)}(s, \boldsymbol{u}) - \hat{\mathbb{Z}}_n^{\star,(m)}(s, \boldsymbol{u}) \right|$$

is the version of $\sup_{s\in[0,1]} \sup_{\boldsymbol{u}\in[0,1]^d} \left| \mathbb{Z}_n^{(m)}(s, \boldsymbol{u}) - \hat{\mathbb{Z}}_n^{(m)}(s, \boldsymbol{u}) \right|$ computed from the sequence $(\xi_n^{(m)}, \boldsymbol{X}_n), (\xi_{n-1}^{(m)}, \boldsymbol{X}_{n-1}), \dots, (\xi_1^{(m)}, \boldsymbol{X}_1)$, which implies that the latter quantity and K_n have the same distribution. Lemma 2 therefore implies that $K_n \overset{\mathrm{P}}{\to} 0$. From the latter fact, Lemma 5 and since $C^{[j]}_n(\boldsymbol{u}) \leq 5$ for all $\boldsymbol{u} \in [0,1]^d$ and $n \geq 1$, we thus obtain that

$$\left(\mathbb{C}_n^\star, \hat{\mathbb{C}}_n^{\star,(1)}, \dots, \hat{\mathbb{C}}_n^{\star,(M)}, \check{\mathbb{C}}_n^{\star,(1)}, \dots, \check{\mathbb{C}}_n^{\star,(M)}\right) \rightsquigarrow \left(\mathbb{C}_C^\star, \mathbb{C}_C^{\star,(1)}, \dots, \mathbb{C}_C^{\star,(M)}, \mathbb{C}_C^{\star,(1)}, \dots, \mathbb{C}_C^{\star,(M)}\right)$$

in $\{\ell^\infty([0,1]^{d+1})\}^{2M+1}$.

From (47), the continuous mapping theorem and the fact that A_n and $A_n^\star$ (defined in (34) and (50), respectively) converge to zero in probability, we have that the weak convergences stated in (48) and (51) occur jointly. This implies that the weak convergence stated in the previous equation occurs jointly with that stated in Theorem 1 (ii). The desired result finally follows from the continuous mapping theorem. ■

Proof of Proposition 5. We first prove (i). Recall that $\boldsymbol{U}_1, \dots, \boldsymbol{U}_n$ is the unobservable sample obtained by applying probability integral transforms to the coordinate sample of $\boldsymbol{X}_1, \dots, \boldsymbol{X}_n$, and, for any $(s, \boldsymbol{u}) \in [0,1]^{d+1}$, let

$$\tilde{\mathbb{D}}_n(s, \boldsymbol{u}) = \sqrt{n}\, \lambda_n(s) \{1 - \lambda_n(s)\} \left[G_{\lfloor ns \rfloor}\{G^{-1}_{\lfloor ns \rfloor}(\boldsymbol{u})\} - G^\star_{n-\lfloor ns \rfloor}\{G^{\star,-1}_{n-\lfloor ns \rfloor}(\boldsymbol{u})\} \right],$$

where, for any $k \in \{1, \dots, n-1\}$,

$$G_k(\boldsymbol{u}) = \frac{1}{k} \sum_{i=1}^k \mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}) \qquad \text{and} \qquad G^\star_{n-k}(\boldsymbol{u}) = \frac{1}{n-k} \sum_{i=k+1}^n \mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}), \quad \boldsymbol{u} \in [0,1]^d,$$

with the convention that $G_0(\boldsymbol{u}) = 0$ and $G_0^\star(\boldsymbol{u}) = 0$, and where, for any $k \in \{1,\dots,n-1\}$ and $\boldsymbol{u} \in [0,1]^d$, $G_k^{-1}(\boldsymbol{u})$ and $G_{n-k}^{\star,-1}(\boldsymbol{u})$ are defined analogously to (38).

Now, $n^{-1/2}\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d}|\mathbb{D}_n(s,\boldsymbol{u}) - \tilde{\mathbb{D}}_n(s,\boldsymbol{u})| \le A_n + B_n$, where

$$A_n = \max_{1\le k\le n} \frac{k(n-k)}{n^2} \sup_{\boldsymbol{u}\in[0,1]^d} \left|C_k(\boldsymbol{u}) - G_k\{G_k^{-1}(\boldsymbol{u})\}\right|$$

and

$$B_n = \max_{1\le k\le n} \frac{k(n-k)}{n^2} \sup_{\boldsymbol{u}\in[0,1]^d} \left|C_{n-k}^\star(\boldsymbol{u}) - G_{n-k}^\star\{G_{n-k}^{\star,-1}(\boldsymbol{u})\}\right|.$$

Then, from (42),

$$A_n \le \max_{1\le k\le n} \frac{k(n-k)}{n^2}\frac{d}{k} \le \frac{d}{n} \qquad \text{and} \qquad B_n \le \max_{1\le k\le n} \frac{k(n-k)}{n^2}\frac{d}{n-k} \le \frac{d}{n},$$

which implies that $n^{-1/2}\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d}|\mathbb{D}_n(s,\boldsymbol{u}) - \tilde{\mathbb{D}}_n(s,\boldsymbol{u})| \le 2d/n$. It follows that, to show (i), we can show the result with $\mathbb{D}_n$ replaced by $\tilde{\mathbb{D}}_n$.

Next, notice that, for any $s \in [0,1]$ and $\boldsymbol{u} \in [0,1]^d$,

$$K_t(s,\boldsymbol{u}) = \begin{cases} s(1-s)C_1(\boldsymbol{u}) - s\{(t-s)C_1(\boldsymbol{u}) + (1-t)C_2(\boldsymbol{u})\} & \text{if } s \le t, \\ (1-s)\{tC_1(\boldsymbol{u}) + (s-t)C_2(\boldsymbol{u})\} - s(1-s)C_2(\boldsymbol{u})\} & \text{if } s > t. \end{cases}$$

Let us therefore split the supremum over s according to the cases $s \in [0,t]$ and $s \in [t,1]$. Then, we write $\sup_{s\in[0,t]}\sup_{\boldsymbol{u}\in[0,1]^d}|n^{-1/2}\tilde{\mathbb{D}}_n(s,\boldsymbol{u}) - K_t(s,\boldsymbol{u})| \le I_n + J_n$, where

$$I_n = \sup_{s\in[0,t]}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\lambda_n(s)\{1-\lambda_n(s)\}G_{\lfloor ns\rfloor}\{G_{\lfloor ns\rfloor}^{-1}(\boldsymbol{u})\} - s(1-s)C_1(\boldsymbol{u})\right|$$

and

$$J_n = \sup_{s\in[0,t]}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\lambda_n(s)\{1-\lambda_n(s)\}G_{n-\lfloor ns\rfloor}^\star\{G_{n-\lfloor ns\rfloor}^{\star,-1}(\boldsymbol{u})\} - s\{(t-s)C_1(\boldsymbol{u}) + (1-t)C_2(\boldsymbol{u})\}\right|.$$

Now,

$$\begin{aligned} I_n &= \sup_{s\in[0,t]}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\lambda_n(s)\{1-\lambda_n(s)\}\left[G_{\lfloor ns\rfloor}\{G_{\lfloor ns\rfloor}^{-1}(\boldsymbol{u})\} - C_1(\boldsymbol{u})\right]\right| + o(1) \\ &\le \sup_{s\in[0,t]}\sup_{\boldsymbol{u}\in[0,1]^d}\left|n^{-1/2}\mathbb{Z}_n\{s, G_{\lfloor ns\rfloor}^{-1}(\boldsymbol{u})\}\right| \\ &\quad + \max_{1\le k\le \lfloor nt\rfloor}\sup_{\boldsymbol{u}\in[0,1]^d}\left|\frac{k(n-k)}{n^2}\left[C_1\{G_k^{-1}(\boldsymbol{u})\} - C_1(\boldsymbol{u})\right]\right| + o(1), \end{aligned}$$

where $\mathbb{Z}_n$ is defined in (9). The first supremum on the right of the last inequality is smaller than $\sup_{s\in[0,t]}\sup_{\boldsymbol{u}\in[0,1]^d}|n^{-1/2}\mathbb{Z}_n(s,\boldsymbol{u})|$ because $G_{\lfloor ns\rfloor}^{-1}(\boldsymbol{u}) \in [0,1]^d$ for all $(s,\boldsymbol{u}) \in [0,1]^{d+1}$. It therefore converges to zero in probability as $\sup_{s\in[0,t]}\sup_{\boldsymbol{u}\in[0,1]^d}|\mathbb{Z}_n(s,\boldsymbol{u})|$ converges in distribution. From (46) and (44), the second term on the right of the previous inequality restricted to $k \geq 3$ is smaller than

$$\max_{3 \leq k \leq \lfloor nt \rfloor} \frac{k}{n} \sum_{j=1}^{d} \sup_{u_j \in [0,1]} \left| G_{k,j}^{-1}(u_j) - u_j \right| \leq \max_{3 \leq k \leq \lfloor nt \rfloor} \frac{k}{n} a_k \times \sum_{j=1}^{d} \max_{3 \leq k \leq \lfloor nt \rfloor} \frac{\sup_{u_j \in [0,1]} |G_{k,j}(u_j) - u_j|}{a_k}$$

$$\leq a_n \times d \max_{1 \leq j \leq d} \sup_{k \geq 3} \frac{\sup_{u_j \in [0,1]} |G_{k,j}(u_j) - u_j|}{a_k},$$

where $a_k = k^{-1/2}(\log\log k)^{1/2}$, $k \geq 3$. By (43), the second term in the product on the right is strictly smaller than ∞ almost surely. Since the first term converges to zero, we obtain that $I_n \overset{\mathrm{P}}{\to} 0$.

Let us now show that $J_n \overset{\mathrm{P}}{\to} 0$. We have $J_n \leq L_n + M_n$, where

$$L_n = \sup_{s \in [0,t]} \sup_{\boldsymbol{u} \in [0,1]^d} \Big| \lambda_n(s) \{1 - \lambda_n(s)\} G^{\star}_{n-\lfloor ns \rfloor} \{ G^{\star,-1}_{n-\lfloor ns \rfloor}(\boldsymbol{u}) \}$$
$$- s \Big[(t-s) C_1 \{ G^{\star,-1}_{n-\lfloor ns \rfloor}(\boldsymbol{u}) \} + (1-t) C_2 \{ G^{\star,-1}_{n-\lfloor ns \rfloor}(\boldsymbol{u}) \} \Big] \Big|$$

and

$$M_n = \sup_{s \in [0,t]} \sup_{\boldsymbol{u} \in [0,1]^d} \left| s \left((t-s) \left[C_1 \{ G^{\star,-1}_{n-\lfloor ns \rfloor}(\boldsymbol{u}) \} - C_1(\boldsymbol{u}) \right] + (1-t) \left[C_2 \{ G^{\star,-1}_{n-\lfloor ns \rfloor}(\boldsymbol{u}) \} - C_2(\boldsymbol{u}) \right] \right) \right|.$$

Now,

$$\begin{aligned} L_n &\leq \sup_{s \in [0,t]} \sup_{\boldsymbol{u} \in [0,1]^d} \left| \lambda_n(s) \{1 - \lambda_n(s)\} G^{\star}_{n-\lfloor ns \rfloor}(\boldsymbol{u}) - s \left[(t-s) C_1(\boldsymbol{u}) + (1-t) C_2(\boldsymbol{u}) \right] \right| \\ &\leq \max_{1 \leq k \leq \lfloor nt \rfloor} \frac{k(n-k)}{n^2} \sup_{\boldsymbol{u} \in [0,1]^d} \left| G^{\star}_{n-k}(\boldsymbol{u}) - \frac{\lfloor nt \rfloor - k}{n-k} C_1(\boldsymbol{u}) - \frac{n - \lfloor nt \rfloor}{n-k} C_2(\boldsymbol{u}) \right| + o(1) \\ &\leq L_{n,1} + L_{n,2} + o(1), \end{aligned}$$

where

$$L_{n,1} = \max_{1 \leq k \leq \lfloor nt \rfloor} \frac{k(\lfloor nt \rfloor - k)}{n^2} \sup_{\boldsymbol{u} \in [0,1]^d} \left| \frac{1}{\lfloor nt \rfloor - k} \sum_{i=k+1}^{\lfloor nt \rfloor} \mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}) - C_1(\boldsymbol{u}) \right|$$
$$\leq n^{-1/2} \sup_{s \in [0,t]} \sup_{\boldsymbol{u} \in [0,1]^d} |\mathbb{Z}_n(t, \boldsymbol{u}) - \mathbb{Z}_n(s, \boldsymbol{u})|$$

and

$$L_{n,2} = \sup_{\boldsymbol{u} \in [0,1]^d} \left| \frac{1}{n - \lfloor nt \rfloor} \sum_{i=\lfloor nt \rfloor + 1}^{n} \mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}) - C_2(\boldsymbol{u}) \right| \times \max_{1 \leq k \leq \lfloor nt \rfloor} \frac{k(n - \lfloor nt \rfloor)}{n^2}.$$

From the continuous mapping theorem, we obtain that $L_{n,1} \overset{\mathrm{P}}{\to} 0$. Also, $\boldsymbol{U}_{\lfloor nt \rfloor + 1}, \ldots, \boldsymbol{U}_n$ being a random sample from C_2, we have, from the Glivenko-Cantelli lemma, that $L_{n,2} \xrightarrow{\text{a.s.}} 0$. Hence, $L_n \overset{\mathrm{P}}{\to} 0$.

Using (46) and (43) as previously, we obtain that $M_n \xrightarrow{\text{a.s.}} 0$, which implies that $J_n \xrightarrow{\text{P}} 0$, and therefore that $\sup_{s\in[0,t]}\sup_{\boldsymbol{u}\in[0,1]^d} |n^{-1/2}\tilde{\mathbb{D}}_n(s,\boldsymbol{u}) - K_t(s,\boldsymbol{u})| \xrightarrow{\text{P}} 0$.

Proceeding similarly, one can show that $\sup_{s\in[t,1]}\sup_{\boldsymbol{u}\in[0,1]^d} |n^{-1/2}\tilde{\mathbb{D}}_n(s,\boldsymbol{u}) - K_t(s,\boldsymbol{u})| \xrightarrow{\text{P}} 0$, which implies that $\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d} |n^{-1/2}\tilde{\mathbb{D}}_n(s,\boldsymbol{u}) - K_t(s,\boldsymbol{u})| \xrightarrow{\text{P}} 0$ and therefore the desired result.

Let us now prove (ii). For any $m \in \{1,\dots,M\}$, let

$$\hat{\mathbb{W}}_n^{(m)}(s,\boldsymbol{u}) = \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} \xi_i^{(m)} \{\mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}) - G_{\lfloor ns \rfloor}(\boldsymbol{u})\}, \qquad (s,\boldsymbol{u}) \in [0,1]^{d+1}$$

and

$$\check{\mathbb{W}}_n^{(m)}(s,\boldsymbol{u}) = \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor ns \rfloor} \xi_i^{(m)} \{\mathbf{1}(\boldsymbol{U}_i \leq \boldsymbol{u}) - G_n(\boldsymbol{u})\}, \qquad (s,\boldsymbol{u}) \in [0,1]^{d+1}.$$

From the proof of Theorem 3 of Holmes et al. (2013), $\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d} |\hat{\mathbb{W}}_n^{(m)}(s,\boldsymbol{u})| = O_{\text{P}}(1)$ and $\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d} |\check{\mathbb{W}}_n^{(m)}(s,\boldsymbol{u})| = O_{\text{P}}(1)$ for all $m \in \{1,\dots,M\}$. Then, from Lemma 1, the fact that

$$\tilde{\hat{\mathbb{Z}}}_n^{(m)}(s,\boldsymbol{u}) = \hat{\mathbb{W}}_n^{(m)}\{s, G_{\lfloor ns \rfloor}^{-1}(\boldsymbol{u})\}, \qquad (s,\boldsymbol{u}) \in [0,1]^{d+1},$$

where $\tilde{\hat{\mathbb{Z}}}_n^{(m)}$ is defined in (39), and the fact that

$$\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d} \left| \hat{\mathbb{W}}_n^{(m)}\{s, G_{\lfloor ns \rfloor}^{-1}(\boldsymbol{u})\} \right| \leq \sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d} \left| \hat{\mathbb{W}}_n^{(m)}(s,\boldsymbol{u}) \right|,$$

we obtain that $\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d} |\hat{\mathbb{Z}}_n^{(m)}(s,\boldsymbol{u})| = O_{\text{P}}(1)$. Hence, $\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d} |\hat{\mathbb{C}}_n^{(m)}(s,\boldsymbol{u})| = O_{\text{P}}(1)$ since $C_n^{[j]}(\boldsymbol{u}) \leq 5$ for all $\boldsymbol{u} \in [0,1]^d$, $j \in \{1,\dots,d\}$ and $n \geq 1$. Similarly, using (40), $\check{\mathbb{W}}_n^{(m)}$ and Lemma 4, we obtain that $\sup_{s\in[0,1]}\sup_{\boldsymbol{u}\in[0,1]^d} |\check{\mathbb{C}}_n^{(m)}(s,\boldsymbol{u})| = O_{\text{P}}(1)$.

Proceeding similarly, it can be verified that the corresponding results for $\hat{\mathbb{C}}_n^{\star,(m)}$ and $\check{\mathbb{C}}_n^{\star,(m)}$ hold, which concludes the proof. ■

References

J. Bai. Weak convergence of the sequential empirical processes of residuals in ARMA models. *Annals of Statistics*, 22(4):2051–2061, 1994.

A. Bücher. A note on nonparametric estimation of bivariate tail dependence. 2012. URL `http://www.ruhr-uni-bochum.de/imperia/md/content/mathematik3/publications/taildep_buecher.pdf`.

A. Bücher and H. Dette. A note on bootstrap approximations for the empirical copula process. *Statistics and Probability Letters*, 80(23–24):1925–1932, 2010.

A. Bücher and M. Ruppert. Consistent testing for a constant copula under strong mixing based on the tapered block multiplier technique. *Journal of Multivariate Analysis*, 116: 208–229, 2013.

A. Bücher and S. Volgushev. Empirical and sequential empirical copula processes under serial dependence. *arXiv:1111.2778*, 2011.

K.L. Chung. An estimate concerning the Kolmogoroff limit distribution. *Transactions of the American Mathematical Society*, 67(1):36–50, 1949.

M. Csörgő and L. Horváth. *Limit theorems in change-point analysis.* Wiley Series in Probability and Statistics. John Wiley & Sons, Chichester, UK, 1997.

P. Deheuvels. La fonction de dépendance empirique et ses propriétés: un test non paramétrique d'indépendance. *Acad. Roy. Belg. Bull. Cl. Sci. 5th Ser.*, 65:274–292, 1979.

P. Deheuvels. A non parametric test for independence. *Publications de l'Institut de Statistique de l'Université de Paris*, 26:29–50, 1981.

H. Dehling, D. Vogel, M. Wendler, and D. Wied. An efficient and robust test for a change-point in correlation. *arXiv:1203.4871*, 2013.

R.M. Dudley and W. Philipp. Invariance principles for sums of Banach space valued random elements and empirical processes. *Probability Theory and Related Fields*, 62: 509–552, 1983.

J.H.J. Einmahl. *Multivariate empirical processes.* Volume 32 of CWI Tract, Amsterdam: Stichting Mathematisch Centrum Centrum voor Wiskunde en Informatica, 1987.

J.H.J. Einmahl and D.M. Mason. Laws of the iterated logarithm in the tails of weighted uniform empirical processes. *Annals of Probability*, 16(1):126–141, 1988.

J.-D. Fermanian, D. Radulovic, and M. Wegkamp. Weak convergence of empirical copula processes. *Bernoulli*, 10(5):847–860, 2004.

P. Gänssler and W. Stute. *Seminar on empirical processes.* DMV Seminar 9. Birkhäuser, Basel, 1987.

A.I. Garralda-Guillem. Structure de dépendance des lois de valeurs extrêmes bivariées. *Comptes rendus de l'Académie des sciences de Paris, Série I Mathématique*, 330:593–596, 2000.

C. Genest and J. Segers. On the covariance of the asymptotic empirical copula process. *Journal of Multivariate Analysis*, 101:1837–1845, 2010.

C. Genest, K. Ghoudi, and L.-P. Rivest. Discussion of "Understanding relationships using copulas", by E. Frees and E. Valdez. *North American Actuarial Journal*, 3:143–149, 1998.

C. Genest, B. Rémillard, and D. Beaudoin. Goodness-of-fit tests for copulas: A review and a power study. *Insurance: Mathematics and Economics*, 44:199–213, 2009.

E. Gombay and L. Horváth. Change-points and bootstrap. *Environmetrics*, 10(6), 1999.

E. Gombay and L. Horváth. Rates of convergence for U-statistic processes and their bootstrapped versions. *Journal of Statistical Planning and Inference*, 102:247–272, 2002.

M. Hofert, I. Kojadinovic, M. Maechler, and J. Yan. *copula: Multivariate dependence with copulas*, 2012. URL `http://CRAN.R-project.org/package=copula`. R package version 0.999-6.

M. Holmes, I. Kojadinovic, and J-F. Quessy. Nonparametric tests for change-point detection à la Gombay and Horváth. *Journal of Multivariate Analysis*, 115:16–32, 2013.

L. Horváth and Q.M. Shao. Limit theorems for permutations of empirical processes with applications to change point analysis. *Stochastic Processes and their Applications*, 117: 1870–1888, 2007.

H. Joe. *Multivariate models and dependence concepts.* Chapman and Hall, London, 1997.

A. Khoudraji. *Contributions à l'étude des copules et à la modélisation des valeurs extrêmes bivariées.* PhD thesis, Université Laval, Québec, Canada, 1995.

J. Kiefer. On large deviations of the empiric d.f. of vector chance variables and a law of the iterated logarithm. *Pacific Journal of Mathematics*, 11(2):649–660, 1961.

I. Kojadinovic and J. Yan. A goodness-of-fit test for multivariate multiparameter copulas based on multiplier central limit theorems. *Statistics and Computing*, 21(1):17–30, 2011.

I. Kojadinovic, J. Segers, and J. Yan. Large-sample tests of extreme-value dependence for multivariate copulas. *The Canadian Journal of Statistics*, 39(4):703–720, 2011.

M.R. Kosorok. *Introduction to empirical processes and semiparametric inference.* Springer, New York, 2008.

E. Liebscher. Construction of asymmetric multivariate copulas. *Journal of Multivariate Analysis*, 99:2234–2250, 2008.

D.M. Mason. Bounds for weighted empirical distribution functions. *Annals of Probability*, 9(5):881–884, 1981.

A.J. McNeil, R. Frey, and P. Embrechts. *Quantitative risk management.* Princeton University Press, New Jersey, 2005.

R.B. Nelsen. *An introduction to copulas.* Springer, New-York, 2006. Second edition.

J.-F. Quessy, M. Saïd, and A.-C. Favre. Multivariate Kendall's tau for change-point detection in copulas. *The Canadian Journal of Statistics*, 2013. URL `doi:10.1002/cjs.11150`.

R Development Core Team. *R: A Language and Environment for Statistical Computing.* R Foundation for Statistical Computing, Vienna, Austria, 2012. URL `http://www.R-project.org`. ISBN 3-900051-07-0.

B. Rémillard. Goodness-of-fit tests for copulas of multivariate time series. *Social Science Research Network*, 1729982:1–32, 2010. URL `http://ssrn.com/abstract=1729982`.

B. Rémillard and O. Scaillet. Testing for equality between two copulas. *Journal of Multivariate Analysis*, 100(3):377–386, 2009.

L. Rüschendorf. Asymptotic distributions of multivariate rank order statistics. *Annals of Statistics*, 4:912–923, 1976.

J. Segers. Asymptotics of empirical copula processes under nonrestrictive smoothness assumptions. *Bernoulli*, 18:764–782, 2012.

G.R. Shorack and J.A. Welner. *Empirical processes with applications to statistics.* Wiley Series in Probability and Mathematical Statistics. John Wiley and Sons, New York, 1986.

A. Sklar. Fonctions de répartition à n dimensions et leurs marges. *Publications de l'Institut de Statistique de l'Université de Paris*, 8:229–231, 1959.

W. Stute. The oscillation behavior of empirical processes: The multivariate case. *Annals of Probability*, 12(2):361–379, 1984.

H. Tsukahara. Semiparametric estimation in copula models. *The Canadian Journal of Statistics*, 33(3):357–375, 2005.

A.W. van der Vaart and J.A. Wellner. *Weak convergence and empirical processes.* Springer, New York, 2000. Second edition.

A.W. van der Vaart and J.A. Wellner. Empirical processes indexed by estimated functions. In *Asymptotics: Particles, Processes and Inverse Problems*, pages 234–252. Institute of Mathematical Statistics, 2007.

M. van Kampen and D. Wied. A nonparametric constancy test for copulas under mixing conditions. *working paper*, 2012. URL `http://www.statistik.tu-dortmund.de/fileadmin/user_upload/Lehrstuehle/Finanzoekonometrie/VanKampenWied_010512.pdf`.

D. Wied, H. Dehling, M. van Kampen, and D. Vogel. A fluctuation test for constant spearman's rho with nuisance-free limit distribution. *arXiv:1206.5070*, 2013.

Table 6: Percentage of rejection of H_0 computed from 1000 samples of size $n \in \{50, 100, 200\}$ such the $\lfloor nt_1 \rfloor$ first observations of each sample are from a d-dimensional c.d.f. with normal copula and $N(0,1)$ margins (that is, from a multivariate standard normal c.d.f.), and the $n - \lfloor nt_1 \rfloor$ last observations are from a d-dimensional c.d.f. with normal copula whose first margin is the $N(\mu, 1)$ and whose d-1 remaining margins are the $N(0,1)$. The bivariate margins of the normal copula have a Kendall's tau of τ.

	$(\mu, t_1) =$		$(0.5, 0.25)$				$(0.5, 0.5)$				$(2, 0.25)$				$(2, 0.5)$			
d	n	τ	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n	$\hat{S}_n$	$\check{S}_n$	S_n^R	κ_n
2	50	0.00	6.7	3.8	9.0	5.3	6.5	3.1	17.7	5.2	4.6	2.4	70.1	3.3	5.1	2.6	98.5	3.9
		0.25	5.6	2.8	8.1	4.6	4.8	3.6	15.0	3.7	6.0	3.1	57.7	4.8	4.7	1.6	98.5	3.9
		0.50	4.4	4.1	9.6	2.5	3.4	2.9	13.0	2.9	18.4	5.1	39.6	14.6	5.7	0.9	99.3	6.4
	100	0.00	5.3	4.0	19.0	4.7	6.3	5.2	30.4	6.4	6.1	4.1	99.0	6.9	5.5	2.5	100.0	5.6
		0.25	5.0	4.4	13.9	5.2	3.7	2.3	24.9	3.6	8.6	4.7	97.5	7.3	4.4	2.2	100.0	5.5
		0.50	4.0	3.8	11.5	3.8	3.1	3.1	22.6	2.1	29.5	13.1	91.6	31.7	12.4	2.0	100.0	18.3
	200	0.00	5.5	5.2	34.1	6.4	3.9	3.4	61.9	5.6	4.3	3.3	100.0	5.2	5.6	4.1	100.0	5.1
		0.25	3.9	3.4	27.6	4.1	4.1	3.5	51.3	5.0	13.6	8.9	100.0	14.6	8.1	4.0	100.0	9.1
		0.50	3.4	3.9	18.9	4.0	2.8	2.9	43.1	3.5	57.8	39.4	100.0	62.2	34.8	8.3	100.0	42.4
3	50	0.00	4.9	1.6	5.0	4.2	4.5	1.7	10.0	4.3	4.8	1.9	36.5	3.7	6.0	2.3	79.6	3.5
		0.25	5.0	2.7	6.9	5.1	5.0	2.6	9.9	4.6	6.9	2.6	24.8	6.6	5.1	2.6	87.4	5.0
		0.50	3.7	2.6	5.5	3.8	3.3	1.3	8.9	4.0	9.4	3.5	17.4	11.6	3.0	0.7	94.2	5.5
	100	0.00	4.5	2.3	11.3	5.2	4.2	2.5	18.6	4.7	4.8	3.0	87.5	5.4	3.6	2.0	99.6	3.2
		0.25	4.9	3.1	10.7	6.0	5.1	3.5	14.8	5.3	6.7	3.8	67.9	8.1	5.9	4.2	99.9	7.3
		0.50	2.8	2.0	7.3	3.8	3.0	2.4	13.7	3.4	16.9	8.6	60.6	22.4	6.0	1.3	100.0	14.9
	200	0.00	3.0	2.4	20.1	4.2	4.5	4.0	37.3	4.8	5.3	3.3	100.0	5.2	4.3	3.2	100.0	5.7
		0.25	4.8	4.1	15.3	5.5	5.6	4.4	30.6	7.3	11.9	8.1	99.2	12.9	7.4	5.2	100.0	8.4
		0.50	4.9	3.7	11.8	5.8	3.8	3.2	24.9	3.5	41.0	30.9	99.2	48.0	23.9	7.8	100.0	32.0